Test Booklet Code

JMD

the second second second

7384287



Do not open this Test Booklet until you are asked to do so.

No.

This Test Booklet contains 20 pages.

Important Instructions :

- 1. The Answer Sheet is inside this Test Booklet. When you are directed to open the Test Booklet, take out the Answer Sheet and fill in the particulars on **Side-1** and **Side-2** carefully with **blue/black** ballpoint pen only.
- The test is of 3 hours duration and Test Booklet contains 180 questions. Each question carries 4 marks. For each correct response, the candidate will get 4 marks. For each incorrect response, one mark will be deducted from the total score. The maximum marks are 720.
- 3. Use Blue/Black Ballpoint Pen only for writing particulars on this page/marking responses.
- 4. Rough work is to be done on the space provided for this purpose in the Test Booklet only.
- 5. On completion of the test, the candidate must hand over the Answer Sheet to the Invigilator before leaving the Room/Hall. The candidates are allowed to take away Test Booklet only with them.
- 6. The CODE for this Test Booklet is **RR**. Make sure that the CODE printed on Side-2 of the Answer Sheet is the same as that on this Test Booklet. In case of discrepancy, the candidate should immediately report the matter to the Invigilator for replacement of both the Test Booklet and the Answer Sheet.
- 7. The candidate should ensure that the Answer Sheet is not folded. Do not make any stray marks on the Answer Sheet. Do not write your Roll No. anywhere else except in the specified space in the Test Booklet/Answer Sheet.
- 8. Use of white fluid for correction is not permissible on the Answer Sheet.
- 9. Each candidate must show on demand his/her Admit Card to the Invigilator.
- 10. No candidate, without special permission of the Superintendent or Invigilator, would leave his/her seat.
- 11. The candidates should not leave the Examination Hall without handing over their Answer Sheet to the Invigilator on duty and sign the Attendance Sheet twice. Cases where a candidate has not signed the Attendance Sheet second time will be deemed not to have handed over the Answer Sheet and dealt with as an unfair means case.
- 12. Use of Electronic/Manual Calculator is prohibited.
- 13. The candidates are governed by all Rules and Regulations of the Board with regard to their conduct in the Examination Hall. All cases of unfair means will be dealt with as per Rules and Regulations of the Board.
- 14. No part of the Test Booklet and Answer Sheet shall be detached under any circumstances.
- 15. The candidates will write the correct Test Booklet Code as given in the Test Booklet/Answer Sheet in the Attendance Sheet.

Name of the Candidate (in Capitals) : Rakshata RAKSHATA TRAD.
Roll Number (in Figures) : 84801.367.
(in Words) : Fight crores forty eight lakhs one thousand three Centre of Examination (in Capitals) : BIRLA SCHOOL KALYAN (W) hundred and sixty-sever
Candidate's Signature : Trabisheita Invigilator's Signature : monthe
Facsimile Signature Stamp of Centre Superintendent :
E3



1.	Which	one	of	the	fo	llowing	ger	nerates	new
	genetic	com	bir	ation	ns	leading	to	variatio	on?
	AT Sex	rual	rep	rodu	cti	ion			

- (2) Nucellar polyembryony
- (3) Vegetative reproduction
- (4) Parthenogenesis
- 2. Match Column-I with Column-II and select the correct option using the codes given below :
- Column-I Column-II a. Pistils fused (i) Gametogenesis together
- b. Formation of (ii) Pistillate gametes
- c. Hyphae of higher (iiii) Syncarpous Ascomycetes
- (iv) Dikaryotic d. Unisexual female flower
 - Codes :

	a	Ъ	с	d	
(1)	(i)	(ii)	(iv)	(iii)	
(2)	(iii)	(i)	(iv)	(ii)	
(3)	(iv)	(iii)	(i)	(ii)	
(4)	(ii)	(i)	(iv)	(iii)	

3. In majority of angiosperms

- (I) reduction division occurs in the megaspore mother cells
- (2) a small central cell is present in the embryo sac
- (3) cgg has a filiform apparatus ×
- (4) there are numerous antipodal cells×
- 4. Pollination in water hyacinth and water lily is brought about by the agency of
 - (1) birds
 - (2) bats
 - (3) water
 - (4) insects or wind
- 5. The ovule of an angiosperm is technically equivalent to
 - (1) megaspore mother cell
 - (2) megaspore
 - (3) megasporangium
 - (4) megasporophyll

JMD/E3

- 6. Taylor conducted the experiments to prove semiconservative .mode of chromosome replication on
 - (1) Drosophila melanogaster
 - (2) E. coli
 - (3) Vinca rosea
 - 141 Vicia faba
- 7. The mechanism that causes a gene to move from one linkage group to another is called
 - AT translocation
 - (2) crossing-over
 - (3) inversion *
 - (4) duplication *
- 8. The equivalent of a structural gene is
 - IT operon
 - (2) recon
 - (3) muton
 - (4) cistron '
- 9. A true breeding plant is
 - (1) near homozygous and produces offspring of its own kind
 - (2) always homozygous recessive in its genetic constitution
 - (3) one that is able to breed on its own
 - (4) produced due to cross-pollination among unrelated plants
- 10. Which of the following rRNAs acts as structural RNA as well as ribozymc in bacteria?
 - (1) 23 S rRNA
 - (2) 5.8 S rRNA
 - (3) 5 S rRNA >
 - 18 S rRNA

- 11. Stirred-tank bioreactors have been designed for
 - (1) availability of oxygen throughout the process
 - 191 ensuring anaerobic conditions in the culture vessel
 - (3) purification of product
 - (4) addition of preservatives to the product

- 12. A forcign DNA and plasmid cut by the same restriction endonuclease can be joined to form a recombinant plasmid using
 - (1) polymerase III
 - (2) ligase
 - (3) Eco RI
 - (4) Taq polymerase

13. Which of the following is not a component of downstream processing?

- (1) Preservation
- (2) Expression
- (3) Separation
- (4) Purification
- 14. Which of the following restriction enzymesproduces blunt ends?
 - (1) Xho I
 - (2) Hind III
 - (3) Sal I
 - (4) Eco RV
- 15. Which kind of therapy was given in 1990 to a four-year-old girl with adenosine deaminase (ADA) deficiency?
 - (1) Immunotherapy
 - (2) Radiation therapy
 - (3) Gene therapy
 - (4) Chemotherapy
- 16. How many hot spots of biodiversity in the world have been identified till date by Norman Myers?

3

- (1) 34
- (2) 43
- (3) 17
- (4) 25

JMD/E3

17. The primary producers of the deep-sea A hydrothermal vent ecosystem are In blue-green algae (2) coral reefs 🛪 ⁽³⁾ green algae (4) chemosynthetic bacteria 18. Which of the following is correct for 4 r-selected species? (1) Small number of progeny with small size (2) Small number of progeny with large size / (3) Large number of progeny with small size (4) Large number of progeny with large size 19. If '+' sign is assigned to beneficial interaction, 1/ '-' sign to detrimental and '0' sign to neutral interaction, then the population interaction represented by '+' '-' refers to (1) commensalism (2) parasitism (3) mutualism (4) amensalism 20. Which of the following is correctly matched? (1) Parthenium hysterophorus-Threat to biodiversity (2) Stratification-Population (3) Aerenchyma-Opuntia (4) Age pyramid-Biome 21. Red List contains data or information on (1) threatened species (2) marine vertebrates only (3) all economically important plants whose products are in (4) plants international trade

[P.T.O.



adel phois-

- **22.** Which one of the following is **wrong** for fungi?
 - (1) They are heterotrophic.
 - (2) They are both unicellular and multicellular.
 - (3) They are eukaryotic.
 - [4] All fungi possess a purely cellulosic cell wall.
- 23. Methanogens belong to
 - (1) Dinoflagellates
 - (2) Slime moulds
 - (3) Eubacteria
 - (A) Archaebacteria
- 24. Select the wrong statement.
 - (1) Diatoms are chief producers in the oceans.
 - (2) Diatoms are microscopic and float passively in water.
 - (3) The walls of diatoms are easily destructible.
 - (4) 'Diatomaceous earth' is formed by the cell walls of diatoms.
- **25.** The label of a herbarium sheet **does not** carry information on
 - (1) local names
 - (2) height of the plant
 - (3) date of collection
 - (4) name of collector
- **26.** Conifers are adapted to tolerate extreme environmental conditions because of
 - It thick cuticle /
 - (2) presence of vessels
 - (3) broad hardy leaves "
 - (4) superficial stomata
- 27. Which one of the following statements is wrong ?
 - (1) Agar-agar is obtained from *Gelidium* and *Gracilaria*.
 - (2) Laminaria and Sargassum arc used as food?
 - (3) Algae increase the level of dissolved oxygen in the immediate environment.
 - Algin is obtained from red algae, and carrageenan from brown algae.

- 28. The term 'polyadelphous' is related to
 - (1) corolla
 - (2) calyx
 - (3) gynoccium
 - (A) and roecium
- **29.** How many plants among *Indigofera*, *Sesbania*, *Salvia*, *Allium*, *Aloe*, mustard, groundnut, radish, gram and turnip have stamens with different lengths in their flowers?
 - (1) Five
 - (2) Six
 - (3) Three
 - (A) Four

30. Radial symmetry is found in the flowers of

- (1) Pisum
- (2) Cassia
- (3) Brassica
- (4) Trifolium
- 31. Free-central placentation is found in
 - (1) Brassica
 - (2) Citrus
 - -(3) Dianthus

4

- (4) Argemone
- 32. Cortex is the region found between
 - (1) endodermis and pith
 - (2) endodermis and vascular bundle
 - (3) epidermis and stele
 - (4) pericycle and endodermis

33. The balloon-shaped structures called tyloses

- (1) are extensions of xylem parenchyma cells into vessels
- (2) are linked to the ascent of sap through xylem vessels
- (3) originate in the lumen of vessels
- (A) characterize the sapwood -

JMD/E3

- 34. A non-proteinaceous enzyme is
 - (1) ligasc
 - (2) deoxyribonuclease
 - (3) lysozyme
 - (4) ribozyme
- 35. Select the mismatch.
 - (1) Protists-Eukaryotes
 - (2) Methanogens-Prokaryotes
 - (3) Gas vacuoles-Green bacteria
 - (4) Large central vacuoles--Animal cells

36. Select the wrong statement.

- (1) Cyanobacteria lack flagellated cells.
 - (Z) Mycoplasma is a wall-less microorganism.
 - (3) Bacterial cell wall is made up of peptidoglycan.
 - (4) Pili and fimbriae are mainly involved in motility of bacterial cells.
- **37.** A cell organelle containing hydrolytic enzymes is
 - (1) ribosome
 - (2) mesosome
 - (3) lysosome
 - (4) microsome
- **38.** During cell growth, DNA synthesis takes place in
 - (1) G₂ phase
 - (2) M phase
 - (3) S phase
 - (4) G₁ phase
- **39.** Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins?
 - (1) Pyruvic acid
 - (2) Acetyl CoA
 - (3) Glucose-6-phosphate
 - (4) Fructosc 1,6-bisphosphate

JMD/E3

- **40.** A few drops of sap were collected by cutting across a plant stem by a suitable method. The sap was tested chemically. Which one of the following test results indicates that it is phloem sap?
 - (1) Low refractive index "
 - (2) Absence of sugar 🛩
 - (3) Acidic r
 - (4) Alkaline r
- **41.** You are given a tissue with its potential for differentiation in an artificial culture. Which of the following pairs of hormones would you add to the medium to secure shoots as well as roots?
 - (I) Auxin and abscisic acid
 - (2) Gibberellin and abscisic acid
 - (3) IAA and gibberellin
 - (4) Auxin and cytokinin
- 42. Phytochrome is a
 - (1) lipoprotein
 - (2) chromoprotein
 - (3) flavoprotein
 - (4) glycoprotein
- 43. Which is essential for the growth of root tip?

(1)	Ca	(2)	Mn
(3)	Zn	 (4)	Fe

- 44. The process which makes major difference between C_3 and C_4 plants is
 - HT photorespiration
 - (2) respiration
 - (3) glycolysis

5

- (4) Calvin cycle
- **45.** Which one of the following statements is **not** correct?
 - .(IT In potato, banana and ginger, the / plantlets arise from the internodes present in the modified stem.
 - (2) Water hyacinth, growing in the standing water, drains oxygen from water that leads to the death of fishes.
 - (3) Offspring produced by the asexual reproduction are called clone.
 - (4) Microscopic, motile asexual reproductive structures are called zoospores.

[P.T.O.

- **46**. The part of nephron involved in active reabsorption of sodium is
 - (1) Bowman's capsule
 - (2) descending limb of Henle's loop
 - (3) distal convoluted tubule
 - 147 proximal convoluted tubule
- **47.** Which of the following is hormonereleasing IUD?
 - (1) Lippes loop
 - (2) Cu7 ×
 - AST LNG-20
 - (4) Multiload 375×
- **48.** Which of the following is **incorrect** regarding vasectomy?
 - (1) Vasa deferentia is cut and tied -
 - (2) Irreversible sterility
 - (3) No sperm occurs in seminal fluid "
 - (14) No sperm occurs in epididymis
- **49.** Embryo with more than 16 blastomeres formed due to *in vitro* fertilization is transferred into
 - (1) fimbriae
 - (2) cervix
 - 131 uterus
 - (4) fallopian tubc
- **50.** Which of the following depicts the **correct** pathway of transport of sperms?
 - Rete testis → Vas deferens → Efferent ductules → Epididymis
 - (2) Efferent ductules → Rete testis → Vas deferens → Epididymis
 - (3) Rete testis \rightarrow Efferent ductules \rightarrow Epididymis \rightarrow Vas deferens
 - (4) Rete testis → Epididymis → Efferent ductules → Vas deferens

(ii) Sperm

(i)

(ii)

Column-II

(i) Embryo formation

(iii) Female external

(iv) Graafian follicle

genitalia

51. Match **Column-I** with **Column-II** and select the correct option using the codes given below :

Column--I

- a. Mons pubis b. Antrum
- c. Trophectoderm

d. Nebenkern

Codes :

	а	b	с	d
(1)	(iii)	(i)	(iv)	(ii)
(2)	(i)	(iv)	(iii)	(ii)

(ii)

(i)

(3) (iii) (iv) (4) (iii) (iv)

JMD/E3

- **52.** Several hormones like hCG, hPL, estrogen, progesterone are produced by
 - (1) fallopian tube
 - (2) pituitary
 - (3) ovary
 - (4) placenta
- **53.** If a colour-blind man marries a woman who is homozygous for normal colour vision, the probability of their son being colour-blind is
 - (1) 0.75 (2) 1
 - (3) 0 (4) 0.5
- 54. Genetic drift operates in
 - (1) non-reproductive population
 - (2) slow reproductive population
 - 135 small isolated population
 - (4) large isolated population
- 55. In Hardy-Weinberg equation, the frequency of heterozygous individual is represented by .
 - (1) pq $f(2) q^2$
 - (3) p^2 (4) 2pq
- **56.** The chronological order of human evolution from early to the recent is
 - (1) Ramapithecus \rightarrow Homo habilis \rightarrow Australopithecus \rightarrow Homo erectus
 - (2) Australopithecus → Homo habilis → Ramapithecus → Homo erectus
 - (3) Australopithecus → Ramapithecus → Homo habilis → Homo erectus
 - .(4) Ramapithecus → Australopithecus → Homo habilis → Homo erectus
- **57.** Which of the following is the **correct** sequence of events in the origin of life?
 - I. Formation of protobionts
 - II. Synthesis of organic monomers
 - III. Synthesis of organic polymers
 - IV. Formation of DNA-based genetic systems
 - (1) II, III, I, IV -
 - 12) II, III, IV, I ~
 - (3) I, II, III, IV ×
 - (4) I, III, II, IV

- 58. A molecule that can act as a genetic material must fulfill the traits given below, except
 - (1) it should be unstable structurally and chemically * -

A DESCRIPTION OF THE OWNER OF THE

- (2) it should provide the scope for slow changes that are required for evolution
- (3) it should be able to express itself in the form of 'Mendelian characters'
- (4) it should be able to generate its replica
- **59.** DNA-dependent RNA polymerase catalyzes transcription on one strend of the DNA which is called the
 - (1) alpha strand
 - (2) antistrand
 - -(3) template strand
 - (4) coding strand
- 60. Interspecific hybridization is the mating of
 - (1) superior males and females of different breeds
 - (2) more closely related individuals within same breed for 4-6 generations
 - (3) animals within same breed without having common ancestors
 - [4] two different related species
- **61.** Which of the following is **correct** regarding AIDS causative agent HIV?
 - (1) HIV is unenveloped retrovirus.
 - (2) HIV does not escape but attacks the acquired immune response.
 - (3) HIV is enveloped virus containing one molecule of single-stranded RNA and one molecule of reverse transcriptase.
 - (4) HIV is enveloped virus that contains two identical molecules of single-stranded RNA and two molecules of reverse transcriptase.
- 62. Among the following edible fishes, which one is a marine fish having rich source of omega-3 fatty acids?
 - (IT Mrigala
 - (2) Mackerel
 - (3) Mystus
 - (4) Mangur

JMD/E3

63. Match **Column-I** with **Column-II** and select the correct option using the codes given below :

	Col	umn-	I		Column-II
a.	Citric	acid		(i)	Trichoderma
b.	Cycle	sporin	А	(ii)	Clostridium
с.	Stati	ns		(iii)	Aspergillus
d.	Buty:	ric acio	1	(iv)	Monascus
Co	des :				
	a	b	с	d	
(1)	(i)	(iv)	(ii)	(iii)	
(2)	(iii)	(iv)	(i)	(ii)	
-13)	(iii)	(i)	(ii)	(iv)	~
(4)	(iii)	(i)	(iv)	(ii)	1

64. Biochemical Oxygen Demand (BOD) may **not** be a good index for pollution for water bodies receiving effluents from

- (1) petroleum industry
- (2) sugar industry
- 137 domestic sewage
- (4) dairy industry
- **65.** The principle of competitive exclusion was stated by
 - (1) MacArthur
 - (2) Verhulst and Pearl
 - (3) C. Darwin
 - (4) G. F. Gause
- **66.** Which of the following National Parks is home to the famous musk deer or hangul?
 - (1) Eaglenest Wildlife Sanctuary, Arunachal Pradesh
 - (2) Dachigam National Park, Jammu & Kashmir
 - (3) Keibul Lamjao National Park, Manipur
 - (4) Bandhavgarh National Park, Madhya & Pradesh

67. A lake which is rich in organic waste may **67** result in

- (1) increased population of fish due to lots of nutrients
- $\mathcal{L}^{(2)}$ mortality of fish due to lack of oxygen
- (3) increased population of aquatic organisms due to minerals
- (4) drying of the lake due to algal bloom

68. The highest DDT concentration in aquatic food chain shall occur in

- (1) crab
- (2) cel
- (3) phytoplankton
- (4) seagull -

organi

[P.T.O.

- **69.** Which of the following sets of diseases is caused by bacteria?
 - (1) Tetanus and mumps
 - (2) Herpes and influenza
 - (3) Cholera and tetanus
 - Typhoid and smallpox :
- 70. Match Column—I with Column—II for housefly classification and select the correct option using the codes given below :

		-		-	
	Col	umn-	-I		Column-II
a.	Fan	nily		(i)	Diptera
Ъ.	Ord	ler ,		-(ii)	Arthropoda
с.	Cla	SS		(iii)	Muscidae
d.	Phy	lum		(iv)	Insecta
Cod	es :				
	a	b	с	d	
1)	(iv)	(iiii)	(ii)	(i)	

- 71. Choose the correct statement.
 - (1) All reptiles have a three-chambered heart.
 - (2) All Pisces have gills covered by an operculum.
 - (3) All mammals are viviparous. *
 - (4) All cyclostomes do not possess jaws and paired fins.
- **72.** Study the four statements (A–D) given below and select the two correct ones out of them :
 - Definition of biological species was given by Ernst Mayr.
 - Photoperiod does not affect reproduction in plants.
 - C. Binomial nomenclature system was given by R. H. Whittaker.
 - D. In unicellular organisms, reproduction is synonymous with growth.

The two correct statements are


- (1) A and D (2) A and B
- (3) B and C (4) C and D
- 73. In male cockroaches, sperms are stored in which part of the reproductive system?(1) Testes
 - (2) Vas deferens
 - 137 Seminal vesicles
 - (4) Mushroom glands
- 74. Smooth muscles are
 - (1) involuntary, cylindrical, striated
 - (2) voluntary, spindle-shaped, uninucleate

and the second second

- (2) involuntary, fusiform, non-striated
- (4) voluntary, multinucleate, cylindrical

JMD/E3

- 75. Oxidative phosphorylation is
 - (1) addition of phosphate group to ATP
 - (2) formation of ATP by energy released from electrons removed during substrate oxidation
 - -(3) formation of ATP by transfer of phosphate group from a substrate to ADP
 - (4) oxidation of phosphate group in ATP
- 76. Which of the following is the least likely to be
- I involved in stabilizing the three-dimensional folding of most proteins?
 - (1) Hydrophobic interaction
 - (2) Ester bonds
 - (3) Hydrogen bonds
 - (4) Electrostatic interaction
- 77. Which of the following describes the given graph correctly?



Reaction \longrightarrow

- Endothermic reaction with energy A in absence of enzyme and B in presence of enzyme
- (2) Exothermic reaction with energy A in absence of enzyme and B in presence of enzyme
- (3) Endothermic reaction with energy A in presence of enzyme and B in absence of enzyme
- (4) Exothermic reaction with energy A in presence of enzyme and B in absence of enzyme
- **78.** When cell has stalled DNA replication fork, which checkpoint should be predominantly activated?
 - (1) M
 - (2) Both G_2/M and M
 - 125 G1/S
 - (4) G_2/M

8

th, S

the second s

5 Con 210 6 0 2

p

					1. 15	5				
					1.2.04	50				
th an giv Co a. Pac	atch the eir char d select ven belo lumn—1 hytene aphase	racteris the co w : I (i)	tic fea rrect o Pairi chro	col ing of	in Colu in Co using umn-	umn- olumi the c -II	n—II odes		acti acti (1) (ð) Nar nur	ne th ive s ivity c Sodin Calci ne th nber
d. Zyg	kinesis otene odes :		Cros Chro	omoso	over tal mes al plate				(1) J2T (3) (4)	Neut Thro Erytl Leuc
(1) (2) (4) (4)	(iv) (iii)	b (iv) (iii) (iv) (iv)	c (iii) (ii) (ii) (ii)	d (i) (i) (i) (iii)	11				on cell (1) (3)	ne a hepa ular Secre Insu
bio	hich I oduction carbona Chole	te?	par	do ncreati nd sec		late ice	the and	87.	ske	letal decro accu
(2) (3) (4)	Insulii Angiot Gastri	n and tensin in and	glucag and e insuli	gon pinepi in	hrine				(3) (4)	infla imm mus high
of (1) (2) (3)	the partia the lun less the less the equal more	igs is han tha han tha to tha	at in at of o t in th	the bl carbon he blo	ood n dioxi od		lveoli		(2) (3) (4)	lacki lacki lacki lacki lacki
	hyper stimu Recep	recepto arized d polarize lus. tors d	rs in luring ed in	the darki respo	humar ness ar nse to	the	come	09.	son	ngs d ne air neve there pres atine there
(3)	pressi	eptors ure.		ond iscles	to ch are		s in rmo-	90.	(4)	there pulli
<pre>(1) (2)</pre>	hypos	isease ecretion secretion	n of a on of n of t	idrena adren hyroic	l gland al gland l gland	ıd I			end	locrin it is thala it se it is it or
JMD /E	23						9			

-55

- e ion responsible for unmasking of ites for myosin for cross-bridge during muscle contraction.
 - (2) Potassium um
 - um (4) Magnesium
- he blood cells, whose reduction in can cause clotting disorder, leading sive loss of blood from the body.
 - rophils 4
 - mbocytes
 - hrocytes
 - ocytes '
- peptide hormone which acts mainly tocytes, adipocytes and enhances glucose uptake and utilization.
 - etin (2) Gastrin
 - (A) Glucagon lin
- rosis, an age-related disease of system, may occur due to
 - eased level of estrogen
 - mulation of uric acid leading to mmation of joints 🛪
 - unc disorder affecting neurocular junction leading to fatigue ¥
 - concentration of Ca⁺⁺ and Na⁺» P-F=1
 - iffers from blood in
 - ng clotting factors
 - ng antibodies
 - ng globulins
 - ng albumins
- o not collapse between breaths and always remains in the lungs which er be expelled because
 - e is a positive intrapleural pressure
 - sure in the lungs is higher than the ospheric pressure
 - e is a negative pressure in the lungs
 - e is a negative intrapleural pressure ng at the lung walls
- terior pituitary gland is not a 'true' e gland because
 - under the regulation of hypoamus
 - cretes enzymes
 - provided with a duct
 - ly stores and releases hormones

0


in

KX

1n


[P.T.O.

91. A parallel-plate capacitor of area A, plate separation d and capacitance C is filled with four dielectric materials having dielectric constants k_1 , k_2 , k_3 and k_4 as shown in the figure below. If a single dielectric material is to be used to have the same capacitance C in this capacitor, then its dielectric constant k is given by



(4)
$$k = \frac{2}{3}(k_1 + k_2 + k_3) + 2k_4$$

92. The potential difference $(V_A - V_B)$ between the points A and B in the given figure is



93. A filament bulb (500 W, 100 V) is to be used in a 230 V main supply. When a resistance *R* is connected in series, it works perfectly and the bulb consumes 500 W. The value of *R* is

(1)	26 Ω	(2)	13 Ω	
(3)	230 Ω	(1)	46 Ω	

94. A long wire carrying a steady current is bent into a circular loop of one turn. The magnetic field at the centre of the loop is B. It is then bent into a circular coil of n turns. The magnetic field at the centre of this coil of n turns will be

(1) 2nB (2) $2n^2B$

(3) nB (4) n^2B

JMD/E3

95. A bar magnet is hung by a thin cotton thread in a uniform horizontal magnetic field and is in equilibrium state. The energy required to rotate it by 60° is W. Now the torque required to kccp the magnet in this new -position is


1)	$\sqrt{3}W$	(2)	2W
	2	(2)	$\sqrt{3}$
3)	W	(4)	$\sqrt{3}W$

96. An electron is moving in a circular path under the influence of a transverse magnetic field of 3.57×10^{-2} T. If the value of e/m is 1.76×10^{11} C/kg, the frequency of revolution of the electron is

(1) 62·8 MHz (2) 6·28 MHz (3) 1 GHz (4) 100 MHz

- **97.** Which of the following combinations should be selected for better tuning of an *L-C-R* circuit used for communication?
 - (1) $R = 15 \Omega$, L = 3.5 H, $C = 30 \mu$ F
 - (2) $R = 25 \Omega$, L = 1.5 H, $C = 45 \mu$ F (3) $R = 20 \Omega$, L = 1.5 H, $C = 35 \mu$ F
 - (4) $R = 25 \Omega$, L = 2.5 H, $C = 45 \mu F$
- **98.** A uniform magnetic field is restricted within a region of radius *r*. The magnetic field changes with time at a rate $\frac{d\vec{B}}{dt}$. Loop 1 of

radius R > r encloses the region r and loop 2 of radius R is outside the region of magnetic field as shown in the figure below. Then the e.m.f. generated is



- (1) $-\frac{d\vec{B}}{dt}\pi R^2$ in loop 1 and zero in loop 2
- (2) $-\frac{dB}{dt}\pi r^2$ in loop 1 and zero in loop 2-
- (3) zero in loop 1 and zero in loop 2

(4)
$$-\frac{dB}{dt}\pi r^2$$
 in loop 1 and

$$\frac{d\vec{B}}{dt}\pi r^2 \text{ in loop } 2$$

50 - 400

- **99.** The potential differences across the resistance, capacitance and inductance are 80 V, 40 V and 100 V respectively in an *L-C-R* circuit. The power factor of this circuit is
 - (1) 0.8 (2) 1.0
 - (3) 0.4 (4) 0.5
- 100. A 100 Ω resistance and a capacitor of 100 Ω reactance are connected in series across a 220 V source. When the capacitor is 50% charged, the peak value of the displacement current is
 - (1) 4.4 A (2) $11\sqrt{2}$ A
 - (3) 2·2 A (4) 11 A
- 101. Two identical glass $(\mu_g = 3/2)$ equiconvex lenses of focal length f each are kept in contact. The space between the two lenses is filled with water $(\mu_w = 4/3)$. The focal length of the combination is
 - (1) 4f/3 (2) 3f/4
 - (3) f/3 (4) f
- 102. An air bubble in a glass slab with refractive index 1.5 (near normal incidence) is 5 cm deep when viewed from one surface and 3 cm deep when viewed from the opposite face. The thickness (in cm) of the slab is
 - (1) 12 (2) 16
 - (3) 8 (4) 10
- 103. The interference pattern is obtained with two coherent light sources of intensity ratio n. In the interference pattern, the ratio



- will be (1) $\frac{\sqrt{n}}{(n+1)^2}$
- (2) $\frac{2\sqrt{n}}{(n+1)^2}$
- (3) $\frac{\sqrt{n}}{n+1}$
- $(4) \quad \frac{2\sqrt{n}}{n+1}$

JMD/E3

- 104. A person can see clearly objects only when they lie between 50 cm and 400 cm from his eyes. In order to increase the maximum distance of distinct vision to infinity, the type and power of the correcting lens, the person has to use, will be
 - (1) concave, -0.2 diopter
 - (2) convex, +0.15 diopter
 - (3) convex, +2.25 diopter
 - (4) concave, -0.25 diopter
- 105. A linear aperture whose width is 0.02 cm is placed immediately in front of a lens of focal length 60 cm. The aperture is illuminated normally by a parallel beam of wavelength 5×10^{-5} cm. The distance of the first dark band of the diffraction pattern from the centre of the screen is
 - (1) 0.20 cm
 - (2) 0.15 cm
 - (3) 0.10 cm
 - (4) 0.25 cm
- 106. Electrons of mass m with de-Broglie wavelength λ fall on the target in an X-ray tube. The cutoff wavelength (λ_0) of the emitted X-ray is

(1)
$$\lambda_0 = \frac{2m^2c^2\lambda^3}{h^2}$$

(2)
$$\lambda_0 = \lambda$$

(3)
$$\lambda_0 = \frac{2mc\lambda^2}{h}$$

(4)
$$\lambda_0 = \frac{2h}{mc}$$

- 107. Photons with energy 5 eV are incident on a cathode C in a photoelectric cell. The maximum energy of emitted photoelectrons is 2 eV. When photons of energy 6 eV are incident on C, no photoelectrons will reach the anode A, if the stopping potential of A relative to C is
 - (1) -1 V
 - (2) -3 V
 - (3) +3 V
 - (4) +4 V

11


[P.T.O.

- from the 3rd orbit to the 2nd orbit, it emits a photon of wavelength λ . When it jumps from the 4th orbit to the 3rd orbit, the corresponding wavelength of the photon will be
 - (1) $\frac{20}{7}\lambda$ (2) $\frac{20}{13}\lambda$ $(3) \quad \frac{16}{25}\lambda \qquad \qquad (4) \quad \frac{9}{16}\lambda$

109. The half-life of a radioactive substance is 30 minutes. The time (in minutes) taken between 40% decay and 85% decay of the same radioactive substance is

(1)	45	(2)	60	
(3)	15	(4)	30	


- 110. For CE transistor amplifier, the audio signal voltage across the collector resistance of $2 k\Omega$ is 4 V. If the current amplification factor of the transistor is 100 and the base resistance is $1 k\Omega$, then the input signal voltage is
 - (2) 15 mV (1) 30 mV
 - (4) 20 mV (3) 10 mV
- 111. The given circuit has two ideal diodes connected as shown in the figure below. The current flowing through the resistance R_1 will be



112. What is the output Y in the following circuit, when all the three inputs A, B, C are first 0 and then 1?

JMD/E3


108. If an electron in a hydrogen atom jumps 113. Planck's constant (h), speed of light in vacuum (c) and Newton's gravitational constant (G) are three fundamental constants. Which of the following combinations of these has the dimension of length?




114. Two cars P and Q start from a point at the same time in a straight line and their positions are represented by $x_P(t) = at + bt^2$ and $x_Q(t) = ft - t^2$. At what time do the cars have the same velocity?

(1) $\frac{a+f}{2(1+b)}$ (2) $\frac{f-a}{2(1+b)}$ (3) $\frac{a-f}{1+b}$ (4) $\frac{a+f}{2(b-1)}$

115. In the given figure, $a = 15 \text{ m/s}^2$ represents the total acceleration of a particle moving in the clockwise direction in a circle of radius R = 2.5 m at a given instant of time. The speed of the particle is



116. A rigid ball of mass m strikes a rigid wall at 60° and gets reflected without loss of speed as shown in the figure below. The value of impulse imparted by the wall on the ball will be



- 117. A bullet of mass 10 g moving horizontally with a velocity of 400 m s⁻¹ strikes a wooden block of mass 2 kg which is suspended by a light inextensible string of length 5 m. As a result, the centre of gravity of the block is found to rise a vertical distance of 10 cm. The speed of the bullet after it emerges out horizontally from the block will be
 - (1) 120 $m s^{-1}$
 - (2) 160 m s⁻¹
 - (3) 100 m s^{-1}
 - (4) 80 m s⁻¹
- 118. Two identical balls A and B having velocities of 0.5 m/s and -0.3 m/s respectively collide elastically in one dimension. The velocities of B and A after the collision respectively will be

(1) -0.3 m/s and 0.5 m/s


- (2) 0.3 m/s and 0.5 m/s
- (3) -0.5 m/s and 0.3 m/s
- (4) 0.5 m/s and -0.3 m/s
- 119. A particle moves from a point $(-2\hat{i} + 5\hat{j})$ to $(4\hat{j} + 3\hat{k})$ when a force of $(4\hat{i} + 3\hat{j})$ N is applied. How much work has been done by the force?
 - (1) 5 J
 - (2) 2 J
 - (3) 8 J
 - (4) 11 J
- 120. Two rotating bodies A and B of masses m and 2m with moments of inertia I_A and $I_B(I_B > I_A)$ have equal kinetic energy of rotation. If L_A and L_B be their angular momenta respectively, then
 - (1) $L_B > L_A$
 - (2) $L_A > L_B$
 - $(3) \quad L_A = \frac{L_B}{2}$
 - (4) $L_A = 2L_B$

JMD/E3

- 121. A solid sphere of mass m and radius R is rotating about its diameter. A solid cylinder of the same mass and same radius is also rotating about its geometrical axis with an angular speed twice that of the sphere. The ratio of their kinetic energies of rotation $(E_{sphere} / E_{cylinder})$ will be
 - (1) 1:4
 - (2) 3:1
 - (3) 2:3
 - (4) 1:5
- 122. A light rod of length l has two masses m_1 and m_2 attached to its two ends. The moment of inertia of the system about an axis perpendicular to the rod and passing through the centre of mass is

(1)
$$(m_1 + m_2)l^2$$
 (2) $\sqrt{m_1m_2}l^2$
(3) $\frac{m_1m_2}{m_1 + m_2}l^2$ (4) $\frac{m_1 + m_2}{m_1m_2}l^2$

123. Starting from the centre of the earth having radius R, the variation of g (acceleration due to gravity) is shown by



124. A satellite of mass m is orbiting the earth (of radius R) at a height h from its surface. The total energy of the satellite in terms of g_0 , the value of acceleration due to gravity at the earth's surface, is

(1)
$$\frac{2mg_0R^2}{R+h}$$

(2)
$$-\frac{2mg_0R^2}{R+h}$$

(3)
$$\frac{mg_0R^2}{2(R+h)}$$

(4)
$$-\frac{mg_0R^2}{2(R+h)}$$

13

[P.T.O.

from $(4 \text{ cm} \times 2 \text{ cm})$ to $(5 \text{ cm} \times 4 \text{ cm})$. If the work done is 3×10^{-4} J, the value of the surface tension of the liquid is

- (1) 0.2 Nm^{-1} (2) 8.0 Nm^{-1} (3) 0.250 Nm^{-1} (4) 0.125 Nm^{-1}
- 126. Three liquids of densities ρ_1 , ρ_2 and ρ_3 (with $\rho_1 > \rho_2 > \rho_3$), having the same value of surface tension T, rise to the same height in three identical capillaries. The angles of contact θ_1 , θ_2 and θ_3 obey
 - (1) $\frac{\pi}{2} < \theta_1 < \theta_2 < \theta_3 < \pi$ (2) $\pi > \theta_1 > \theta_2 > \theta_3 > \frac{\pi}{2}$ (3) $\frac{\pi}{2} > \theta_1 > \theta_2 > \theta_3 \ge 0$ (4) $0 \le \theta_1 < \theta_2 < \theta_3 < \frac{\pi}{2}$
- 127. Two identical bodies are made of a material for which the heat capacity increases with temperature. One of these is at 100 °C, while the other one is at 0 °C. If the two bodies are brought into contact, then, assuming no heat loss, the final common temperature is
 - (1) less than 50 °C but greater than 0 °C
 - (2) 0°C
 - (3) 50 °C
 - (4) more than 50 °C
- 128. A body cools from a temperature 3T to 2T in 10 minutes. The room temperature is T. Assume that Newton's law of cooling is applicable. The temperature of the body at the end of next 10 minutes will be
 - (1) (2) T(4) $\frac{3}{2}T$ (3) $\frac{7}{1}T$
- 129. One mole of an ideal monatomic gas undergoes a process described by the equation PV^3 = constant. The heat capacity of the gas during this process is

(1) $2R$	(2)	R
(3) $\frac{3}{2}R$	(4)	5 K

JMD/E3

125. A rectangular film of liquid is extended 130. The temperature inside a refrigerator is t_2 °C and the room temperature is t_1 °C. The amount of heat delivered to the room for each joule of electrical energy consumed ideally will be

(1)
$$\frac{t_2 + 273}{t_1 - t_2}$$
 (2) $\frac{t_1 + t_2}{t_1 + 273}$
(3) $\frac{t_1}{t_1 - t_2}$ (4) $\frac{t_1 + 273}{t_1 - t_2}$


- 131. A given sample of an ideal gas occupies a volume V at a pressure P and absolute temperature T. The mass of each molecule of the gas is m. Which of the following gives the density of the gas?
 - (2) mkT(1) P/(kTV)

(3) P/(kT)(4) Pm/(kT)

- 132. A body of mass m is attached to the lower end of a spring whose upper end is fixed. The spring has negligible mass. When the mass m is slightly pulled down and released, it oscillates with a time period of $3 ext{ s. }$ When the mass m is increased by 1 kg, the time period of oscillations becomes 5 s. The value of m in kg is
 - (2) $\frac{9}{16}$ (4) $\frac{4}{2}$ (3)
- 133. The second overtone of an open organ pipe has the same frequency as the first overtone of a closed pipe L metre long. The length of the open pipe will be
 - (1) $\frac{L}{2}$ (2) 4L
 - (4) 2L (3) L
- 134. Three sound waves of equal amplitudes have frequencies (n-1), n, (n+1). They superimpose to give beats. The number of beats produced per second will be
 - (1) 3 (2) 2 (3) 1 (4) 4
- 135. An electric dipole is placed at an angle of 30° with an electric field intensity 2×10^5 N/C. It experiences a torque equal to 4 N m. The charge on the dipole, if the dipole length is 2 cm, is
 - (1) 5 mC
 - (2) 7 µC
 - (3) 8 mC
 - (4) 2 mC

14

PALIT I K W 14 Mai (1- (1-) 3 136. Hot concentrated sulphuric acid is a moderately strong oxidizing agent. Which of spin complexes of
142. Jahn-Teller effect is not observed in high e 0 r the following reactions does not show F UN d4 oxidizing behaviour? $(2) d^{9}$ (1) $C + 2H_2SO_4 \rightarrow CO_2 + 2SO_2 + 2H_2O$ (x(3) d7 $(4) d^8$ (2) $CaF_2 + H_2SO_4 \rightarrow CaSO_4 + 2HF$ (3) $Cu + 2H_2SO_4 \rightarrow CuSO_4 + SO_2 + 2H_2O_4$ 143. Which of the following can be used as (4) $3S + 2H_2SO_4 \rightarrow 3SO_2 + 2H_2O$ the halide component for Friedel-Crafts 137. Which of the following pairs of d-orbitals will reaction? have electron density along the axes? H3 HSCI IT Chloroethene (H d 2, d 2-12 (2) $d_{xy}, d_{x^2-y^2}$ (2) Isopropyl chloride (3) d_{2}, d_{xz} (3) Chlorobenzene (4) d_{xx}, d_{yz} (4) Bromobenzene 138. The correct geometry and hybridization for XeF₄ are 144. In which of the following molecules, all 7 (1) planar triangle, sp^3d^3 atoms are coplanar? Lops (2) square planar, sp^3d^2 (3) octahedral, sp^3d^2 (4) trigonal bipyramidal, sp^3d . U. 139. Among the following, which one is a wrong (3) statement? (1) SeF_4 and CH_4 have same shape. (2) I_3^+ has bent geometry. (3) PH₅ and BiCl₅ do not exist. $\mathcal{Y} p\pi$ - $d\pi$ bonds are present in SO₂. 145. Which one of the following structures represents nylon 6,6 polymer? 140. The correct increasing order of trans-effect 4 of the following species is (1) $Br^- > CN^- > NH_3 > C_6H_5^-$ (2) $CN^- > Br^- > C_6H_5^- > NH_3$ /(3) NH₃ > CN⁻ > Br⁻ > C₆H₅ $[4]^{-}CN^{-} > C_{6}H_{5}^{-} > Br^{-} > NH_{3}$ 141. Which one of the following statements related to lanthanons is incorrect? (1) All the lanthanons are much more reactive than aluminium. (2) Ce(+4) solutions are widely used as oxidizing agent in volumetric analysis. Is Europium shows +2 oxidation state. (4) The basicity decreases as the ionic radius decreases from Pr to Lu. JMD/E3 15 [P.T.O. S TE NA AI Li Be OC No Ma Als:




the electron density is maximum on

- (1) 2 and 4
- (2) 2 and 5
- (3) 2 and 3
- ×(4) 3 and 4

147. Which of the following compounds shall not produce propene by reaction with HBr followed by elimination or direct only elimination reaction? 6-6-6

- (2) H_3C —C— CH_2Br
- H₂C CH₂ + Mas Br_L (3)(4) H₃C-C-CH₂OH


148. Which one of the following nitro-compounds does not react with nitrous acid?




JMD/E3

149. The central dogma of molecular genetics states that the genetic information flows from


- (1) DNA \rightarrow RNA \rightarrow Proteins
 - (2) DNA \rightarrow RNA \rightarrow Carbohydrates
 - (3) Amino acids \rightarrow Proteins \rightarrow DNA
 - (4) DNA → Carbohydrates → Proteins
- 150. The correct corresponding order of names of four aldoses with configuration given below




151. In the given reaction



the product P is



 $u = c - N \theta_{L}$ $u = c - N \theta_{L}$



	internet internet
0 H. H. O. H. O.	cits coon n= 3pp title
158. Which one of the following compounds	163. How many electrons can fit in the orbital for which $n = 3$ and $l = 1$?
shows the presence of intramolecular hydrogen bond?	(1) 10 (2) 14
(1) Cellulose 0	(3) 2. 147 6
(2) Concentrated acetic acid x	164. For a sample of perfect gas when its pressure
$CH_3 COOM$ (2) Concentrated acetic acid x (3) H_2O_2	is changed isothermally from p_i to p_f , the
(4) HCN	entropy change is given by (p_c)
159. The molar conductivity of a $0.5 \text{ mol}/\text{dm}^3$	(1) $\Delta S = nRT \ln\left(\frac{p_f}{p_i}\right)$
solution of AgNO3 with electrolytic	
conductivity of 5.76×10^{-3} S cm ⁻¹ at	(2) $\Delta S = RT \ln \left(\frac{p_i}{p_f} \right)$
298 K is (1) $a \cos^2 (\cos^2)$	
(1) $0.086 \text{ S cm}^2 / \text{mol}$	(3) $\Delta S = nR \ln \left(\frac{P_f}{P_i}\right)$
(2) $28 \cdot 8 \text{ S cm}^2 / \text{mol}$	
(3) $2.88 \text{ S cm}^2 / \text{mol}$	(4) $\Delta S = nR \ln\left(\frac{p_i}{p_f}\right)$
(4) 11.52 S cm^2 /mol	165. The van't Hoff factor (i) for a dilute aqueous
160. The decomposition of phosphine (PH3) on	solution of the strong electrolyte barium
tungsten at low pressure is a first-order	hydroxide is
reaction. It is because the	(1) 2 (2) 3 (3) 0 (4) 1
(1) rate is independent of the surface coverage	166. The percentage of pyridine (C_5H_5N) that
(2) rate of decomposition is very slow	forms pyridinium ion $(C_5H_5N^+H)$ in a 0.10 M
(3) rate is proportional to the surface	aqueous pyridine solution $(K_b$ for
coverage	$C_5H_5N = 1.7 \times 10^{-9}$) is
(4) rate is inversely proportional to the surface coverage	C ₅ H ₅ N = 1.7×10^{-9}) is (1) 0.77% C ₅ H ₅ N (2) 1.6% 0.1° $1^{1/2}$ (3) 0.0060% 1 (4) 0.013% 7 167. In calcium fluoride, having the fluorite
161. The coagulation values in millimoles per litre	(3) 0.0060% M
of the electrolytes used for the coagulation of	(4) 0.013% X
As_2S_3 are given below :	167. In calcium fluoride, having the fluorite
I. $(NaCl) = 52$, II. $(BaCl_2) = 0.69$,	structure, the coordination numbers for calcium ion (Ca ²⁺) and fluoride ion (F ⁻) are
$A_{32}S_3$ III. (MgSO ₄) = 0.22 The correct order of their coagulating power is	
power is	121 4 and 8
(1) (1) (1) (1) (2) (1) (1) (1)	, 043) 4 and 2
$(3) I > II > III \qquad (4) II > I > III$	AT 6 and 6
162. During the electrolysis of molten sodium	168. If the \vec{E}_{cell} for a given reaction has a negative
chloride, the time required to produce 0.10 mol of chlorine gas using a current of	f value, which of the following gives the correct relationships for the values of ΔQ
3 amperes is	and Keq?
(1) 220 minutes	(1) $\Delta G^{\circ} < 0; K_{eq} > 1$
(2) 330 minutes	(2) $\Delta G^{\circ} < 0; K_{eq} < 1$
(3) 55 minutes	(3) $\Delta G^{\circ} > 0; K_{eq} < 1$
(4) 110 minutes	(4) $\Delta G^{\circ} > 0; K_{cq} > 1$
IMD/E3 +2 -2 -2	18
JMD/E3 taFL ZCAF AT 6 CAFL AT 6 F 8 6	S CONSTR CAFI) LT
AT 6 Carz &	S C C C C C C F
r y Ca	F23 B Ca

169. Which one of the following is incorrect for 175. Zinc can be coated on iron to produce ideal solution? galvanized iron but the reverse is not (1) $\Delta P = P_{obs} - P_{calculated by Raoult's law} = 0$ possible. It is because (1) zinc has lower negative electrode (2) $\Delta G_{\text{mix}} = 0$ potential than iron (3) $\Delta H_{\rm mix} = 0$ he (2) zinc has higher negative electrode (4) $\Delta U_{\rm mix} = 0$ potential than iron 170. The solubility of AgCl (s) with solubility (3) zinc is lighter than iron product 1.6×10^{-10} in 0.1 M NaCl solution (4) zinc has lower melting point than iron would be 176. The suspension of slaked lime in water is (1) 1.6 × 10⁻¹¹ M known as (1) milk of lime (2) zero (2) aqueous solution of slaked lime (3) $1.26 \times 10^{-5} M$ (3) limewater (4) $1.6 \times 10^{-9} M$ (4) quicklime 15 171. Suppose the elements X and Y combine to 177. The hybridizations of atomic orbitals of form two compounds XY_2 and X_3Y_2 . When nitrogen in NO2, NO3 and NH4 respectively 0.1 mole of XY₂ weighs 10 g and 0.05 mole are of X_3Y_2 weighs 9 g, the atomic weights of X and Y are (1) sp, sp^2 and sp^3 11 (x) 20, 30 (2) 30, 20 (2) sp^2 , sp and sp^3 1 ×3~12 (4) 60, 40 T (3) 40, 30 (3) sp, sp^3 and sp^2 (4) sp^2 , sp^3 and sp172. The number of electrons delivered at the cathode during electrolysis by a current of l ampere in 60 seconds is (charge on 178. Which of the following fluoro-compounds is $electron = 1.60 \times 10^{-19} C$ most likely to behave as a Lewis base? (1) CF₄ (1) 3.75×10^{20} (2) 7.48×10^{23} (2) SiF_4 (3) BF3 (4) PF3 (3) 6×10^{23} (4) 6×10^{20} 179. Which of the following pairs of ions is isoelectronic and isostructural? 173. Boric acid is an acid because its molecule (1) SO_3^{2-} , NO_3^{-} (2) ClO_3^{-} , SO_3^{2-} (1) accepts OH⁻ from water releasing proton (3) CO_3^{2-} , NO_3^{-} (4) ClO_3^{-} , CO_3^{2-} (2) combines with proton from water molecule (3) contains replaceable H⁺ ion 180. In context with beryllium, which one of the following statements is incorrect? (4) gives up a proton (1) Its salts rarely hydrolyze. 174. AlF₃ is soluble in HF only in presence of KF. (2) Its hydride is electron-deficient and It is due to the formation of polymeric. (1) AlH₃ (2) $K[AIF_2H]$ (3) It is rendered passive by nitric acid. (3) $K_3[A1F_3H_3]$ (4) $K_3[A1F_6]$ Li Br (4) It forms Be C. Na Ma Acid Ar. JMD/E3 Be [P.T.O. log. Ma Ma AN SI PS Rh Sn 5. H NO3 G 134 10g.

Test Booklet Code

JMD

This Test Booklet contains 20 pages.

A States

AA

3

3

Do not open this Test Booklet until you are asked to do so.

Important Instructions :

- 1. The Answer Sheet is inside this Test Booklet. When you are directed to open the Test Booklet, take out the Answer Sheet and fill in the particulars on Side-1 and Side-2 carefully with **blue/black** ballpoint pen only.
- The test is of 3 hours duration and Test Booklet contains 180 questions. Each question carries 4 marks. For each correct response, the candidate will get 4 marks. For each incorrect response, one mark will be deducted from the total score. The maximum marks are 720.
- 3. Use Blue/Black Ballpoint Pen only for writing particulars on this page/marking responses.
- 4. Rough work is to be done on the space provided for this purpose in the Test Booklet only.
- 5. On completion of the test, the candidate must hand over the Answer Sheet to the Invigilator before leaving the Room/Hall. The candidates are allowed to take away Test Booklet only with them.
- 6. The CODE for this Test Booklet is **AA**. Make sure that the CODE printed on **Side-2** of the Answer Sheet is the same as that on this Test Booklet. In case of discrepancy, the candidate should immediately report the matter to the Invigilator for replacement of both the Test Booklet and the Answer Sheet.
- 7. The candidate should ensure that the Answer Sheet is not folded. Do not make any stray marks on the Answer Sheet. Do not write your Roll No. anywhere else except in the specified space in the Test Booklet/Answer Sheet.
- 8. Use of white fluid for correction is not permissible on the Answer Sheet.
- 9. Each candidate must show on demand his/her Admit Card to the Invigilator.
- 10. No candidate, without special permission of the Superintendent or Invigilator, would leave his/her seat.
- 11. The candidates should not leave the Examination Hall without handing over their Answer Sheet to the Invigilator on duty and sign the Attendance Sheet twice. Cases where a candidate has not signed the Attendance Sheet second time will be deemed not to have handed over the Answer Sheet and dealt with as an unfair means case.
- 12. Use of Electronic/Manual Calculator is prohibited.
- 13. The candidates are governed by all Rules and Regulations of the Board with regard to their conduct in the Examination Hall, All cases of unfair means will be dealt with as per Rules and Regulations of the Board.
- 14. No part of the Test Booklet and Answer Sheet shall be detached under any circumstances.
- 15. The candidates will write the correct Test Booklet Code as given in the Test Booklet/Answer Sheet in the Attendance Sheet.

Name of the Candida	te (in Capitals) :_	RAAFIAH	IZHAR		
Roll Number (in Figu	ares) :8	1418841	18		
(in Wor	rds) : <u>£1441</u>	Due Four Du	e Eight Ei	ght Four	r Oue Sutdexlu
Centre of Examinatio	n (in Capitals) : _	DEIHI POLICE	PUBLIC	SCHOL	B-4 Enclare Dell
Candidate's Signature	: Kaufiah	Yahan Invigilato	or's Signature : St	dut	B-4 Enclave Del
Facsimile Signature S Centre Superintender	-			•	90 - 10
E1		1. J			
	ά.	S			

Planck's constant (h), speed of light in vacuum (c) and Newton's gravitational three fundamental constant (G) are constants. Which of the following combinations of these has the dimension of length?

 $\frac{Gc}{h^{3/2}}$

(1)
$$\frac{\sqrt{hG}}{c^{3/2}}$$
 (2)
(3) $\sqrt{\frac{hc}{G}}$ (4)

a

CA

à

2. Two cars P and Q start from a point at the same time in a straight line and their positions are represented by $x_P(t) = at + bt^2$ and $x_O(t) = ft - t^2$. At what time do the cars

have the same velocity?

(1)
$$\frac{a-f}{1+b}$$
 (2) $\frac{a+f}{2(b-1)}$
(3) $\frac{a+f}{2(1+b)}$ (4) $\frac{f-a}{2(1+b)}$

In the given figure, $a = 15 \text{ m/s}^2$ represents the total acceleration of a particle moving in the clockwise direction in a circle of radius R = 2.5 m at a given instant of time. The speed of the particle is

$$4.5 \text{ m/s}$$
 (2) 5.0 m/s
5.7 m/s (4) 6.2 m/s

A. A rigid ball of mass m strikes a rigid wall at 60° and gets reflected without loss of speed as shown in the figure below. The value of impulse imparted by the wall on the ball will be

> (2) 2mV $(4)^{-} \frac{mV}{mV}$

> > (? ??

2 LINE

$$\frac{\sqrt{k^{2}}}{\sqrt{m}}$$

(1) (3)

(5.) A bullet of mass 10 g moving horizonta with a velocity of 400 m s^{-1} strikes a wood block of mass 2 kg which is suspended by light inextensible string of length 5 m. As result, the centre of gravity of the block found to rise a vertical distance of 10 cr The speed of the bullet after it emerges or horizontally from the block will be

- (1) 100 m s^{-1}
- (2) 80 m s⁻¹
- (3) 120 m s^{-1}
- (4) 160 $m s^{-1}$
- . Two identical balls A and B having velocities of 0.5 m/s and -0.3 m/s respectively collide elastically in one dimension. The velocities of B and A after the collision respectively will be
 - (1) -0.5 m/s and 0.3 m/s
 - (2) 0.5 m/s and -0.3 m/s
 - (3) -0.3 m/s and 0.5 m/s
 - (4) 0.3 m/s and 0.5 m/s
- *A*. A particle moves from a point $(-2\hat{i} + 5\hat{j})$ to $(4\hat{j}+3\hat{k})$ when a force of $(4\hat{i}+3\hat{j})$ N is applied. How much work has been done by the force?

(8.) Two rotating bodies A and B of masses mand 2m with moments of inertia I_A and $I_B(I_B > I_A)$ have equal kinetic energy of rotation. If L_A and L_B be their angular momenta respectively, then $(1) \quad L_A = \frac{L_B}{2}$ $(2) \quad L_A = 2L_B$ $(3) \quad L_B > L_A$ $(4) \quad L_A > L_B$

KE2 JW2




A solid sphere of mass m and radius R is rotating about its diameter. A solid cylinder of the same mass and same radius is also rotating about its geometrical axis with an angular speed twice that of the sphere. The ratio of their kinetic energies of rotation $(E_{\text{sphere}} / E_{\text{cylinder}})$ will be

- (1) 2:3
- (2) 1:5
- (3) 1:4
- (4) 3:1

10. A light rod of length l has two masses m_1 and m_2 attached to its two ends. The moment of inertia of the system about an axis perpendicular to the rod and passing through the centre of mass is

- (1) $\frac{m_1m_2}{m_1 + m_2} l^2$ (2) $\frac{m_1 + m_2}{m_1m_2} l^2$ (3) $(m_1 + m_2) l^2$ (4) $\sqrt{m_1m_2} l^2$
- 11. Starting from the centre of the earth having radius R, the variation of g (acceleration due to gravity) is shown by



KETIN

- 13. A rectangular film of liquid is extended from $(4 \text{ cm} \times 2 \text{ cm})$ to $(5 \text{ cm} \times 4 \text{ cm})$. If the work done is 3×10^{-4} J, the value of the surface tension of the liquid is
 - (1) 0.250 Nm^{-1} (2) 0.125 Nm^{-1}
 - (3) 0.2 Nm^{-1} (4) 8.0 Nm^{-1}
- 14. Three liquids of densities ρ_1 , ρ_2 and ρ_3 (with $\rho_1 > \rho_2 > \rho_3$), having the same value of surface tension *T*, rise to the same height in three identical capillaries. The angles of contact θ_1 , θ_2 and θ_3 obey

(1)
$$\frac{\pi}{2} > \theta_1 > \theta_2 > \theta_3 \ge 0$$

(2) $0 \le \theta_1 < \theta_2 < \theta_3 < \frac{\pi}{2}$
(3) $\frac{\pi}{2} < \theta_1 < \theta_2 < \theta_3 < \pi$
(4) $\pi > \theta_1 > \theta_2 > \theta_3 > \frac{\pi}{2}$
(5) $\frac{\pi}{2}$

15. Two identical bodies are made of a material for which the heat capacity increases with temperature. One of these is at 100 °C, while the other one is at 0 °C. If the two bodies are brought into contact, then, assuming no heat loss, the final common temperature is (1) 50 °C

- (2) more than 50 °C
- (3) less than 50 °C but greater than 0 °C
- _(4) 0 °C

A body cools from a temperature 3T to 2Tin 10 minutes. The room temperature is T. Assume that Newton's law of cooling is applicable. The temperature of the body at the end of next 10 minutes will be

(1) $\frac{7}{4}T$ (2) $\frac{3}{2}T$ (3) $\frac{4}{3}T$ (4) T

One mole of an ideal monatomic gas undergoes a process described by the equation PV^3 = constant. The heat capacity of the gas during this process is

(2) $\frac{5}{2}R$ (1) $\frac{3}{2}R$ (3) 2R(4) RT = (n-2T)10 K 2 T

(18.) The temperature inside a refrigerator is t_2 °C and the room temperature is t_1 °C. The amount of heat delivered to the room for each joule of electrical energy consumed ideally will be

t_1	(2) $\frac{t_1 + 273}{2}$		
(1) $\frac{t_1}{t_1 - t_2}$	$t_1 - t_2$		
3) $\frac{t_2 + 273}{4}$	(4) $\frac{t_1 + t_2}{1 - 272}$		
$t_1 - t_2$	$t_1 + 273$		

- 19. A given sample of an ideal gas occupies a volume V at a pressure P and absolute temperature T. The mass of each molecule of the gas is m. Which of the following gives the density of the gas?
 - (1) P/(kT)(2) Pm/(kT)(3) P / (kTV) (4) mkT
- **20.** A body of mass m is attached to the lower end of a spring whose upper end is fixed. The spring has negligible mass. When the mass m is slightly pulled down and released, it oscillates with a time period of 3 s. When the mass m is increased by 1 kg, the time period of oscillations becomes 5 s. The value of m in kg is

 $(1) \frac{3}{2}$ 10 16

L

2

(1) 1

(3) 3

(2) 3 $\frac{3}{16}$ (4) $\frac{3}{16}$ (5) The second overtone of an open organ

pipe has the same frequency as the first overtone of a closed pipe L metre long. The length of the open pipe will be (2) 2L

nor shout A

m(1) L+(3)


- (4) 4L
- 2) Three sound waves of equal amplitudes have frequencies (n-1), n, (n+1). They superimpose to give beats. The number of beats produced per second will be

(2) 4

- (4) 2
- An electric dipole is placed at an angle of 30° with an electric field intensity 2×10^5 N/C. It experiences a torque equal to 4 Nm. The charge on the dipole, if the dipole length is 2 cm, is
 - (1) 8 mC
 - (2) 2 mC
 - (3) 5 mC
 - (4) 7 µC

JMD/E1

24. A parallel-plate capacitor of area A, plate separation d and capacitance C is filled with four dielectric materials having dielectric constants k_1 , k_2 , k_3 and k_4 as shown in the figure below. If a single dielectric material is to be used to have the same capacitance C in this capacitor, then its dielectric constant kis given by



72 mg

32. The potential differences across the resistance, capacitance and inductance are 80 V, 40 V and 100 V respectively in an L-C-R circuit. The power factor of this circuit is

(1) 0.4 (2) 0.5

(3) 0.8 (4) 1.0

- **33.** A 100 Ω resistance and a capacitor of 100 Ω reactance are connected in series across a 220 V source. When the capacitor is 50% charged, the peak value of the displacement current is
 - (1) 2·2 A (2) 11 A
 - (4) $11\sqrt{2}$ A (3) 4·4 A
- **34.** Two identical glass $(\mu_g = 3/2)$ equiconvex lenses of focal length f each are kept in contact. The space between the two lenses is filled with water ($\mu_w = 4/3$). The focal length of the combination is

(1)	f/3	(2) f
(-)	J / -	

- (3) 4f/3(4) 3f/4
- 35. An air bubble in a glass slab with refractive index 1.5 (near normal incidence) is 5 cm deep when viewed from one surface and 3 cm deep when viewed from the opposite face. The thickness (in cm) of the slab is

(1)	8	See.	(2)	10
(3)	12		(4)	16

36. The interference pattern is obtained with two coherent light sources of intensity ratio n. In the interference pattern, the ratio

$$\frac{I_{\max} - I_{\min}}{I_{\max} + I_{\min}}$$

will be

(1) $\frac{\sqrt{n}}{n+1}$ (2) $\frac{2\sqrt{n}}{n+1}$ (3) $\frac{\sqrt{n}}{(n+1)^2}$ (4) $\frac{2\sqrt{n}}{(n+1)^2}$

$$\int (L-C)^2 + R^2$$

[P.T.O.

3. A bar magnet is hung by a thin cotton thread in a uniform horizontal magnetic field and is in equilibrium state. The energy required to rotate it by 60° is W. Now the torque required to keep the magnet in this new position is

W (2) $\sqrt{3}W$ (1) $\sqrt{3}$ (3) $\frac{\sqrt{3}W}{2}$ (4) $\frac{2W}{\sqrt{2}}$

29. An electron is moving in a circular path under the influence of a transverse magnetic field of 3.57×10^{-2} T. If the value of e/m is 1.76×10^{11} C/kg, the frequency of revolution of the electron is

(1)	1 GHz	(2)	100	MHz


(3) 62·8 MHz (4) 6.28 MHz

Which of the following combinations should be selected for better tuning of an L-C-R circuit used for communication?

(1) $R = 20 \Omega, L = 1.5 H, C = 35 \mu F$

- (2) $R = 25 \Omega, L = 2.5 H, C = 45 \mu F$
- (3) $R = 15 \Omega, L = 3.5 H, C = 30 \mu F$
- (4) $R = 25 \Omega, L = 1.5 H, C = 45 \mu F$

A uniform magnetic field is restricted within a region of radius r. The magnetic field changes with time at a rate $\frac{d\vec{B}}{dt}$. Loop 1 of radius R > r encloses the region r and loop 2 of radius R is outside the region of magnetic field as shown in the figure below. Then the e.m.f. generated is




- (1) zero in loop 1 and zero in loop 2 (2) $-\frac{d\vec{B}}{dt}\pi r^2$ in loop 1 and
 - $-\frac{dB}{dt}\pi r^2$ in loop 2
- (3) $-\frac{d\vec{B}}{dt}\pi R^2$ in loop 1 and zero in loop 2 (4) $-\frac{dB}{dt}\pi r^2$ in loop 1 and zero in loop 2

JMD/E1

- 37. A person can see clearly objects only when they lie between 50 cm and 400 cm from his eyes. In order to increase the maximum distance of distinct vision to infinity, the type and power of the correcting lens, the person has to use, will be
 - (1) convex, +2.25 diopter
 - (2) concave, -0.25 diopter
 - (3) concave, -0.2 diopter
 - (4) convex, +0.15 diopter
- **38.** A linear aperture whose width is 0.02 cm is placed immediately in front of a lens of focal length 60 cm. The aperture is illuminated normally by a parallel beam of wavelength 5×10^{-5} cm. The distance of the first dark band of the diffraction pattern from the centre of the screen is
 - (1) 0.10 cm
 - (2) 0.25 cm
 - (3) 0.20 cm
 - (4) 0.15 cm

. Electrons of mass *m* with de-Broglie wavelength λ fall on the target in an X-ray tube. The cutoff wavelength (λ_0) of the emitted X-ray is



Photons with energy 5 eV are incident on a cathode C in a photoelectric cell. The maximum energy of emitted photoelectrons is 2 eV. When photons of energy 6 eV are incident on C, no photoelectrons will reach the anode A, if the stopping potential of A relative to C is

- (1) +3 V
 - (2) +4 V
 - (3) -1 V
 - (4) -3 V

JMD/E1


41. If an electron in a hydrogen atom jumps from the 3rd orbit to the 2nd orbit, it emits a photon of wavelength λ . When it jumps from the 4th orbit to the 3rd orbit, the corresponding wavelength of the photon will be

(1)	$\frac{16}{25}\lambda$	2	(2)	$\frac{9}{16}\lambda$
(3)	$\frac{20}{7}\lambda$		(4)	$\frac{20}{13}\lambda$


42. The half-life of a radioactive substance is 30 minutes. The time (in minutes) taken between 40% decay and 85% decay of the same radioactive substance is

(1)	15	(2)	30
(3)	45	(4)	60

- 43. For CE transistor amplifier, the audio signal voltage across the collector resistance of $2 k\Omega$ is 4 V. If the current amplification factor of the transistor is 100 and the base resistance is $1 k\Omega$, then the input signal voltage is
 - (1) 10 mV (2) 20 mV
 - (3) 30 mV (4) 15 mV
- 44. The given circuit has two ideal diodes connected as shown in the figure below. The current flowing through the resistance R_1 5218 15 will be



45. What is the output Y in the following circuit, when all the three inputs A, B, C are first 0 and then 1?



\sim			1
46	Which one of the following compounds	51.	
\sim	shows the presence of intramolecular hydrogen bond?		wh
~	H_2O_2		(1)
	(2) HCN	()	(3)
	(3) Cellulose	192.	Fo
	(4) Concentrated acetic acid		en
47	The molar conductivity of a $0.5 \text{ mol}/\text{dm}^3$		(1)
<i>ч1</i> .	solution of $AgNO_3$ with electrolytic		
	conductivity of $5.76 \times 10^{-3} \text{ S cm}^{-1}$ at		(2)
	298 K is		(2)
	(1) 2.88 S cm^2 /mol		(3)
	(2) 11.52 S cm^2 /mol	i	ý
	(3) 0.086 S cm^2 /mol		(4)
	(4) 28.8 S cm^2 /mol		(+)
	(1) 28.8 5 cm / mor	(53).	Th
(48).	The decomposition of phosphine (PH_3) on		sol
V	tungsten at low pressure is a first-order reaction. It is because the		hy
			(1)
C	(1) rate is proportional to the surface coverage	54.	(3)
	(2) rate is inversely proportional to the	54.	for
· *	surface coverage		aq
	(3) rate is independent of the surface		C ₅
	coverage		(1)
	(4) rate of decomposition is very slow.		(2)
49.	The coagulation values in millimoles per litre of the electrolytes used for the coagulation of	1. A.	(3)
	As $_2S_3$ are given below :		(4)
	I. $(NaCl) = 52$, II. $(BaCl_2) = 0.69$,	55.	,In str
	III. $(MgSO_4) = 0.22$		cal
	The correct order of their coagulating		(1)
	power is		(2)
	$(1) I > II > III \qquad (2) II > I > III (2) II > I > III $		(3)
	$(3) III > II > I \qquad (4) III > I > II$		(4)
50.	During the electrolysis of molten sodium	56.	If t
	chloride, the time required to produce 0.10 mol of chlorine gas using a current of		val
	3 amperes is		an
	(1) 55 minutes		(1)
	(2) 110 minutes		(2)
	(3) 220 minutes		(3)
	(4) 330 minutes		(4)
	177.1	_	
JMD	/EI	7	

How many electrons can fit in the orbital for which n = 3 and l = 1?

(1) 2 (2) 6 (3) 10 (4) 14

2. For a sample of perfect gas when its pressure is changed isothermally from p_i to p_f , the entropy change is given by

(1) $\Delta S = nR \ln \left(\frac{p_f}{p_i}\right)$ $(2) \Delta S = nR \ln \left(\frac{p_i}{p_f}\right)$ (3) $\Delta S = nRT \ln \left(\frac{p_f}{p_i'}\right)$ $(3f' \Delta S = nRT \ln \left(\frac{p_f}{p_i'}\right)$ $(4) \Delta S = RT \ln \left(\frac{p_i}{p_f}\right)$

53. The van't Hoff factor (i) for a dilute aqueous solution of the strong electrolyte barium hydroxide is

- (1) 0 (2) 1
- (3) 2 (4) 3

54. The percentage of pyridine (C_5H_5N) that forms pyridinium ion ($C_5H_5N^+H$) in a 0.10 \dot{M} aqueous pyridine solution (K_b for $C_5H_5N = 1.7 \times 10^{-9}$) is

- (1) 0.0060%
- (2) 0.013%
- (3) 0.77%
- (4) 1.6%
- **55.** In calcium fluoride, having the fluorite structure, the coordination numbers for calcium ion (Ca²⁺) and fluoride ion (F⁻) are
 - (1) 4 and 2
 - (2) 6 and 6
 - (3) 8 and 4
 - (4) 4 and 8
- **56.** If the E_{cell}° for a given reaction has a negative value, which of the following gives the **correct** relationships for the values of ΔG° and K_{eq} ?
 - (1) $\Delta G^{\circ} > 0$; $K_{eq} < 1$


(2) $\Delta G^{\circ} > 0$; $K_{eq} > 1$

(3) $\Delta G^{\circ} < 0$; $K_{eq} > 1$


- (4) $\Delta G^{\circ} < 0$; $K_{eq} < 1$
- [P.T.O.

63. Zinc can be coated on iron to produce Which one of the following is **incorrect** for galvanized iron but the reverse is not ideal solution? possible. It is because (1) $\Delta H_{\rm mix} = 0$ (1) zinc is lighter than iron (2) $\Delta U_{\rm mix} = 0$ (2) zinc has lower melting point than iron (3) $\Delta P = P_{obs} - P_{calculated by Raoult's law} = 0$ (3) zinc has lower negative electrode (4) $\Delta G_{\text{mix}} = 0$ potential than iron 58. The solubility of AgCl (s) with solubility (4) zinc has higher negative electrode product 1.6×10^{-10} in 0.1 M NaCl solution potential than iron would be 64. The suspension of slaked lime in water is (1) $1.26 \times 10^{-5} M$ known as Mh (1) limewater (2) $1.6 \times 10^{-9} M$ (2) quicklime (3) $1.6 \times 10^{-11} M$ (3) milk of lime (4) zero (4) aqueous solution of slaked lime 65. The hybridizations of atomic orbitals of **59.** Suppose the elements X and Y combine to form two compounds XY₂ and X₃Y₂. When nitrogen in NO₂⁺, NO₃⁻ and NH₄⁺ respectively 0.1 mole of XY₂ weighs 10 g and 0.05 mole (1) sp, sp^3 and sp^2 5 + 18 + (2) sp^2 , sp^3 and sp 6 284 (3) (3) sp, sp^2 and sp^3 6 214 5 are of X₃Y₂ weighs 9 g, the atomic weights of X and Y are (2) sp^2 , sp^3 and sp(2) sp, sp^2 and sp^3 (1) 40, 30 (2) 60, 40 (3) 20, 30 (4) 30, 20 (4) sp^2 , sp and sp^3 60. The number of electrons delivered at the cathode during electrolysis by a current of 1 ampere in 60 seconds is (charge on 66. Which of the following fluoro-compounds is most likely to behave as a Lewis base? electron = 1.60×10^{-19} C) (1) BF₃ (2) PF₃ (1) 6×10^{23} (2) 6×10^{20} (3) CF₄ (4) SiF_4 (3) 3.75×10^{20} (4) 7.48×10^{23} 67. Which of the following pairs of ions is isoelectronic and isostructural? 61. Boric acid is an acid because its molecule $(1) CO_3^{2-}, NO_3^{-}$ (2) ClO_3^{-}, CO_3^{2-} (1) contains replaceable H⁺ ion (3) SO_3^{2-} , NO_3^{-} (4) CIO_3^{-} , SO_3^{2-} (2) gives up a proton (3) accepts OH⁻ from water releasing proton 68. In context with beryllium, which one of the (4) combines with proton from water following statements is incorrect? molecule (1) It is rendered passive by nitric acid. **62.** AlF₃ is soluble in HF only in presence of KF. (2) It forms Be₂C. It is due to the formation of (3) Its salts rarely hydrolyze. (1) $K_3[AIF_3H_3]$ (2) $K_3[AIF_6]$ (4) Its hydride is electron-deficient and (4) $K[A1F_3H]$ (3) AlH₃ polymeric. WA+8+2 78 JMD/E1 8

69. Hot concentrated sulphuric acid is a 75. Jahn-Teller effect is not observed in high moderately strong oxidizing agent. Which of spin complexes of the following reactions does not show (1) d^7 (2) d^8 oxidizing behaviour? (1) $Cu + 2H_2SO_4 \rightarrow CuSO_4 + SO_2 + 2H_2O$ (4) d^9 (3) d^4 (2) $3S + 2H_2SO_4 \rightarrow 3SO_2 + 2H_2O_4$ (3) $C + 2H_2SO_4 \rightarrow CO_2 + 2SO_2 + 2H_2O_3$ 76. Which of the following can be used as (4) $CaF_2 + H_2SO_4 \rightarrow CaSO_4 + 2HF$ the halide component for Friedel-Crafts 70. Which of the following pairs of d-orbitals will reaction? have electron density along the axes? Chlorobenzene (1) d_{z^2}, d_{xz} (2) d_{xz}, d_{yz} Bromobenzene (3) d_{2}, d_{2}, d_{2} (3) Chloroethene (4) $d_{xy}, d_{x^2-y^2}$ (4) Isopropyl chloride 71, The correct geometry and hybridization for XeF₄ are (77. In which of the following molecules, all (1) octahedral, sp^3d^2 atoms are coplanar? (2) trigonal bipyramidal, $sp^3d \neq$ 63 (3) planar triangle, sp^3d^3 (1) 302 M square planar, $sp^{3}d^{2}$ 72. Among the following, which one is a wrong statement? (1) PH₅ and BiCl₅ do not exist. (3)(2) $p\pi$ - $d\pi$ bonds are present in SO₂. (8) SeF₄ and CH₄ have same shape. (4) I₃⁺ has bent geometry. 78. Which one of the following structures represents nylon 6,6 polymer? 73. The correct increasing order of trans-effect of the following species is (1) $NH_3 > CN^- > Br^- > C_6H_5^-$ (2) $CN^- > C_6H_5^- > Br^- > NH_3$ (3) $Br^- > CN^- > NH_3 > C_6H_5^-$ (4) $CN^- > Br^- > C_6H_5^- > NH_3$ 74. Which one of the following statements related to lanthanons is incorrect? (1) Europium shows +2 oxidation state. (2) The basicity decreases as the ionic radius decreases from Pr to Lu. (4) $\begin{pmatrix} H_2 & H_2 \\ C + C & C \\ H_2 & H \\ H_2 & H \end{pmatrix}$ (3) All the lanthanons are much more reactive than aluminium. (4) Ce(+4) solutions are widely used as oxidizing agent in volumetric analysis. [P.T.O. JMD/E1



the product P is



CHO

-OH H

ĊH₂OH

JMD/E1

1

(4)


H₃C


H₃C

NO2

CH3 HNO2

(3) H₃C-






JMD/E1

	\sim
03. A non-proteinaceous enzyme is	A few drops of sap were collected by cutting across a plant stem by a suitable method.
(1) lysozyme	The sap was tested chemically. Which one of
(2) ribozyme	the following test results indicates that it is
(3) ligase	phloem sap?
(4) deoxyribonuclease	(1) Acidic
	(2) Alkaline (3) Low refractive index
104. Select the mismatch.	(4) Absence of sugar
(1) Gas vacuoles—Green bacteria	
(2) Large central vacuoles—Animal cells	(110.) You are given a tissue with its potential for differentiation in an artificial culture. Which
(3) Protists—Eukaryotes	of the following pairs of hormones would you
(4) Methanogens—Prokaryotes	add to the medium to secure shoots as well as roots?
105. Select the wrong statement.	(1) IAA and gibberellin
(1) Bacterial cell wall is made up o	of (2) Auxin and cytokinin
peptidoglycan.	(3) Auxin and abscisic acid
(2) Pili and fimbriae are mainly involved in	n (4) Gibberellin and abscisic acid
motility of bacterial cells.	111. Phytochrome is a
(3) Cyanobacteria lack flagellated cells.	(1) flavoprotein
(4) Mycoplasma is a wall-les	s (2) glycoprotein
microorganism.	(3) lipoprotein
106. A cell organelle containing hydrolyti	c (4) chromoprotein
enzymes is	112. Which is essential for the growth of root tip?
HT lysosome	(1) Zn (2) Fe
(2) microsome	(3) Ca (4) Mn
(3) ribosome	113. The process which makes major difference
(4) mesosome	between C_3 and C_4 plants is
-0-	(1) glycolysis
107 During cell growth, DNA synthesis take	s. (2) Calvin cycle
	(3) photorespiration
(1) S phase	(4) respiration
(2) G ₁ phase	114. Which one of the following statements is not
(3) G ₂ phase	correct?
(4) M phase	(1) Offspring produced by the asexual reproduction are called clone.
108. Which of the following biomolecules i common to respiration-mediated breakdow.	(2) meroscopie, motile ascalar reproductive
of fats, carbohydrates and proteins?	structures are cancu zoospores.
(1) Glucose-6-phosphate	(3) In potato, banana and ginger, the plantlets arise from the internodes
(2) Fructose 1,6-bisphosphate	present in the modified stem.
	(4) Water hyacinth, growing in the standing
(3) Pyruvic acid	water, drains oxygen from water that
(4) Acetyl CoA	leads to the death of fishes.
JMD/E1	13 [P.T.O.

 115. Which one of the following generates new genetic combinations leading to variation? (1) Vegetative reproduction (2) Parthenogenesis (3) Sexual reproduction (4) Nucellar polyembryony 	 120. Taylor conducted the experiments to prove semiconservative mode of chromosome replication on (1) Vinca rosea (2) Vicia faba (3) Drosophila melanogaster
116. Match Column-I with Column-II and select the correct option using the codes given below :	(4) E. coli (121) The mechanism that causes a gene to move from one linkage group to another is called
Column—IColumn—IIa. Pistils fused together(i) Gametogenesis togetherb. Formation of gametes(ii) Pistillate somycetesc. Hyphae of higher Ascomycetes(iii) Syncarpous togetherd. Unisexual female(iv) Dikaryotic	 (1) inversion (2) duplication (3) translocation (3) translocation (4) crossing-over (1) The equivalent of a structural gene is (1) muton
flower Codes :	(2) cistron (3) operon (4) recon
 a b c d y i (1) (iv) (iii) (i) (ii) y (2) (ii) (i) (iv) (iii) y (3) (i) (ii) (iv) (iii) (4) (iii) (i) (iv) (ii) 117. In majority of angiosperms (1) egg has a filiform apparatus (2) there are numerous antipodal cells (3) reduction division occurs in the megaspore mother cells (4) a small central cell is present in the embryo sac 118. Pollination in water hyacinth and water lily is brought about by the agency of (1) water (2) insects or wind (3) birds (4) bats 	 (4) recon 123. A true breeding plant is one that is able to breed on its own produced due to cross-pollination among unrelated plants near homozygous and produces offspring of its own kind (4) always homozygous recessive in its genetic constitution (4) always homozygous recessive in its genetic constitution (4) Which of the following rRNAs acts as structural RNA as well as ribozyme in bacteria? (1) 5 S rRNA (2) 18 S rRNA (3) 23 S rRNA (4) 5.8 S rRNA 125. Stirred-tank bioreactors have been designed for
 119. The ovule of an angiosperm is technically equivalent to (1) megasporangium (2) megasporophyll (3) megaspore mother cell (4) megaspore 	 (1) purification of product (2) addition of preservatives to the product (3) availability of oxygen throughout the process (4) ensuring anaerobic conditions in the culture vessel
JMD /E1 1	.4

Q

0			
128.	A foreign DNA and plasmid cut by the same restriction endonuclease can be joined to form a recombinant plasmid using	131.	The primary producers of the deep-sea hydrothermal vent ecosystem are
	·		(1) green algae
	(1) Eco RI		(2) chemosynthetic bacteria
	(2) Taq polymerase		[3] blue-green algae
	(3) polymerase III		(4) coral reefs
	(4) ligase	-	
127.	Which of the following is not a component of downstream processing?	132.	Which of the following is correct for <i>r</i> -selected species?
\sim			(1) Large number of progeny with small size
	(1) Separation		(2) Large number of progeny with large size
100	(2) Purification		(3) Small number of progeny with small size
((3) Preservation		(4) Small number of progeny with large size
A	(4) Expression		
128.	Which of the following restriction enzymes produces blunt ends?	133.	If '+' sign is assigned to beneficial interaction, '-' sign to detrimental and '0' sign to neutral interaction, then the population interaction represented by '+' '-' refers to
\bigcirc	(1) Sal I		(1) mutualism
	(2) Eco RV		
	(3) Xho I	8 10	(2) amensalism
	(4) Hind III		(3) commensalism
129.	Which kind of therapy was given in 1990 to a four-year-old girl with adenosine deaminase (ADA) deficiency?	(Jan)	Which of the following is correctly matched?
	HT Gene therapy		(1) Aerenchyma—Opuntia
	(2) Chemotherapy		(2) Age pyramid—Biome
	(3) Immunotherapy		(3) Parthenium hysterophorus—Threat to biodiversity
	(4) Radiation therapy		(4) Stratification—Population
130.	How many hot spots of biodiversity in the world have been identified till date by Norman Myers?	135.	Red List contains data or information on
	20.50 20.005		(1) all economically important plants
	 (1) 17 - (2) 25 		(2) plants whose products are in international trade
-	(3) 34		(3) threatened species
	(4) 43		(4) marine vertebrates only
ner		-	
JMD	/E1 1	5	[P.T.O.



16. Match the stages of meiosis in Column-I to | 151. Name the ion responsible for unmasking of their characteristic features in Column-II and select the correct option using the codes given below :

 ColumnI (Paring of homologous chromosomes b. Metaphase I (Paring of homologous chromosomes do stimulation of chiasmata equatorial plate Codes: Codes: Codes: Codes: Codes: Codes: Codes: (a) b c d d (b) (ii) (ii) (ii) (ii) (ii) (ii) (ii) (given below :	(1) Calcium (2) Magnesium
 b. Metaphase I (ii) Terminalization of chiasmata c. Diakinesis (iii) Crossing-over takes place d. Zygoten (iv) Chromosomes align at equatorial plate Codes : a b c / d (i) (ii) (ii) (ii) (2) (i) (iv) (iii) (ii) (3) (ii) (iv) (iii) (ii) (4) (iv) (iii) (ii) (5) (Colecystokinin and secretin (4) Insulin and glucagon (1) Angiotensin and epinephrine (2) Gastrin and insulin (3) Crobecystokinin and secretin (4) Insulin and glucagon (3) East han that in the blood (2) more than that in the blood (3) less than that of carbon dioxide (4) Ises than that of carbon dioxide (4) Ises than that of carbon dioxide (5) Reaves disease is couseles are thermoreceptors. (3) Photoreceptors in the human eye ard cholarized during darkness and become hyperolarized in response to the light stimulus. (4) Receptors do not produce grade optentials. (5) Graves' disease is caused due to (1) hyposecretion of adrenal gland (2) hypersecretion of adrenal gland (3) hyposecretion of adrenal gland (4) hypersecretion of adrenal gland 	Column—I Column—II	(3) Sodium (4) Potassium
 a b c d b d c d c d d d iv (ii) (ii) (ii) d d iv (iii) (ii) (ii) d d v (iii) (iii) (ii) d d v (iii) (ii) (ii) d v (iii) (ii) (ii) d v (iii) (ii) (ii) d d v (iii) (ii) (ii) d v (iii) (ii) (iii) d d v (iii) (iii) (iii) (iii) d d v (iii) (ii	 b. Metaphase I (ii) Terminalization of chiasmata c. Diakinesis (iii) Crossing-over takes place d. Zygotene (iv) Chromosomes align at equatorial plate 	number can cause clotting disorder, leading to excessive loss of blood from the body. (1) Erythrocytes (2) Leucocytes (3) Neutrophils
 (2) hypersecretion of thyroid gland (3) hyposecretion of adrenal gland (4) hypersecretion of adrenal gland (5) hypersecretion of adrenal gland (4) it secretes enzymes 	 a b c d (1) (iii) (iv) (ii) (ii) (i) (2) (i) (iv) (ii) (iii) (iii) (3) (ii) (iv) (iii) (i) (4) (iv) (iii) (ii) (i) (4) (iv) (iii) (ii) (i) (5) (147. Which hormones do stimulate the production of pancreatic juice and bicarbonate? (1) Angiotensin and epinephrine (2) Gastrin and insulin (3) Cholecystokinin and secretin (4) Insulin and glucagon 148. The partial pressure of oxygen in the alveoli of the lungs is (1) equal to that in the blood (2) more than that in the blood (3) less than that in the blood (4) less than that of carbon dioxide (4) Schoecytors respond to changes in pressure. (2) Meissner's corpuscles are thermoreceptors. (3) Photoreceptors in the human eye are depolarized during darkness and become hyperpolarized in response to the light stimulus. (4) Receptors do not produce graded potentials. (150. Graves' disease is caused due to 	 153. Name a peptide hormone which acts mainly on hepatocytes, adipocytes and enhances cellular glucose uptake and utilization. (1) Insulin (2) Glucagon (3) Secretin (4) Gastrin 154. Osteoporosis, an age-related disease of skeletal system, may occur due to (1) immune disorder affecting neuromuscular junction leading to fatigue (2) high concentration of Ca⁺⁺ and Na⁺ (3) decreased level of estrogen (4) accumulation of uric acid leading to inflammation of joints 155. Serum differs from blood in (1) lacking globulins (2) lacking clotting factors (4) lacking antibodies 156. Lungs do not collapse between breaths and some air always remains in the lungs which can never be expelled because (1) there is a negative intrapleural pressure pulling at the lung walls (3) there is a positive intrapleural pressure (4) pressure in the lungs is higher than the atmospheric pressure 157. The posterior pituitary gland is not a 'true' endocrine gland because (1) it's provided with a duct
	(2) hypersecretion of thyroid gland(3) hyposecretion of adrenal gland(4) hypersecretion of adrenal gland	 (3) it is under the regulation of hypo- thalamus (4) it secretes enzymes

[P.T.O.

active sites for myosin for cross-bridge

activity during muscle contraction.

\bigcap	
(158.) The part of nephron involved in active reabsorption of sodium is	164. Several hormones like hCG, hPL, estrogen, progesterone are produced by
(1) distal convoluted tubule	(1) ovary
(2) proximal convoluted tubule	(2) placenta
(3) Bowman's capsule(4) descending limb of Henle's loop	(3) fallopian tube
	(4) pituitary
(159) Which of the following is hormone- releasing IUD?	(4) pituliary
LI LNG-20	165, If a colour-blind man marries a woman who
(2) Multiload 375	is homozygous for normal colour vision, the
(3) Lippes loop	probability of their son being colour-blind is
(4) Cu7	(1) 0 (2) 0·5
160. Which of the following is incorrect regarding vasectomy?	$(3) 0.75 \qquad (4) 1 \qquad \neq \checkmark 1 \rightarrow$
(1) No sperm occurs in seminal fluid	166. Genetic drift operates in
(2) No sperm occurs in epididymis	(1) small isolated population $+$ /
(3) Vasa deferentia is cut and tied(4) Irreversible sterility	(2) large isolated population
161. Embryo with more than 16 blastomeres	(3) non-reproductive population
formed due to <i>in vitro</i> fertilization is	(4) slow reproductive population
transferred into	\sim
(1) uterus	(167) In Hardy-Weinberg equation, the frequency
<(2) fallopian tube	of heterozygous individual is represented by
(3) fimbriae(4) cervix	(1) p^2 $(1) 2pq$
162 . Which of the following depicts the correct	(3) pq (4) q^2
pathway of transport of sperms?	
	168. The chronological order of human evolution
Epididymis \rightarrow Vas deferens	from early to the recent is
(2) Rete testis → Epididymis → Efferent ductules → Vas deferens	(1) Australopithecus \rightarrow Ramapithecus \rightarrow
(3) Rete testis \rightarrow Vas deferens \rightarrow Efferent	Homo habilis → Homo erectus
ductules \rightarrow Epididymis	(2) Ramapithecus \rightarrow Australopithecus \rightarrow
(4) Efferent ductules \rightarrow Rete testis \rightarrow Vas	Homo habilis \rightarrow Homo erectus
deferens \rightarrow Epididymis 163. Match Column–II with Column–II and	(3) Ramapithecus → Homo habilis → Australopithecus → Homo erectus
select the correct option using the codes	(4) Australopithecus \rightarrow Homo habilis \rightarrow
given below :	Ramapithecus \rightarrow Homo erectus
Column—I Column—II	
a. Mons pubis 🔨 (i) Embryo formation	169. Which of the following is the correct sequence of events in the origin of life?
b. Antrum (ii) Sperm	
c. Trophectoderm (iii) Female external	I. Formation of protobionts
d. Nebenkern (iv) Graafian follicle	II. Synthesis of organic monomers
	· III. Synthesis of organic polymers
Codes :	IV. Formation of DNA-based genetic systems
$\begin{array}{cccc} \mathbf{a} & \mathbf{b} & \mathbf{c} & \mathbf{d} \\ (1) & (\mathbf{i}\mathbf{i}\mathbf{i})^{*} & (\mathbf{i}\mathbf{r}\mathbf{i}) & (\mathbf{i}\mathbf{i}) & (\mathbf{i}\mathbf{i}) \end{array}$	(1) I, II, III, IV
(1) (iii) (iv) (ii) (i) (2) (iii) (iv) (i) (ii) \cdot	(2) I, III, II, IV
(3) (iii) (i) (iv) (ii) (ii)	(3) II, III, I, IV
(4) (i) (iv) (iii) (ii)	(4) II, III, IV, I
JMD /E1 13	8
	-

STREET.

۲.

ý.

Test Booklet Code

JMI ai Mata Di

7539134



This Test Booklet contains 20 pages.

No.


Do not open this Test Booklet until you are asked to do so.

Important Instructions:

- 1. The Answer Sheet is inside this Test Booklet. When you are directed to open the Test Booklet, take out the Answer Sheet and fill in the particulars on Side-1 and Side-2 carefully with blue/black ballpoint pen only.
- The test is of 3 hours duration and Test Booklet contains 180 questions. Each question carries 4 marks. For each correct response, the candidate will get 4 marks. For each incorrect response, one mark will be deducted from the total score. The maximum marks are 720.
- 3. Use Blue/Black Ballpoint Pen only for writing particulars on this page/marking responses.
- 4. Rough work is to be done on the space provided for this purpose in the Test Booklet only.
- 5. On completion of the test, the candidate must hand over the Answer Sheet to the Invigilator before leaving the Room/Hall. The candidates are allowed to take away Test Booklet only with them.
- 6. The CODE for this Test Booklet is **XX**. Make sure that the CODE printed on **Side-2** of the Answer Sheet is the same as that on this Test Booklet. In case of discrepancy, the candidate should immediately report the matter to the Invigilator for replacement of both the Test Booklet and the Answer Sheet.
- 7. The candidate should ensure that the Answer Sheet is not folded. Do not make any stray marks on the Answer Sheet. Do not write your Roll No. anywhere else except in the specified space in the Test Booklet/Answer Sheet.
- 8. Use of white fluid for correction is not permissible on the Answer Sheet.
- 9. Each candidate must show on demand his/her Admit Card to the Invigilator.
- 10. No candidate, without special permission of the Superintendent or Invigilator, would leave his/her seat.
- 11. The candidates should not leave the Examination Hall without handing over their Answer Sheet to the Invigilator on duty and sign the Attendance Sheet twice. Cases where a candidate has not signed the Attendance Sheet second time will be deemed not to have handed over the Answer Sheet and dealt with as an unfair means case.
- 12. Use of Electronic/Manual Calculator is prohibited.
- 13. The candidates are governed by all Rules and Regulations of the Board with regard to their conduct in the Examination Hall. All cases of unfair means will be dealt with as per Rules and Regulations of the Board.
- 14. No part of the Test Booklet and Answer Sheet shall be detached under any circumstances.
- 15. The candidates will write the correct Test Booklet Code as given in the Test Booklet/Answer Sheet in the Attendance Sheet.

Name of the Candidate (in Capitals) : MEMUL AUTT
Roll Number (in Figures) : 672.00254
(in Words) : Eight Gore Seeruh Juo hakh & Two hundredfifty
Centre of Examination (in Capitals) : KV No 2, MBK, D-dun
Candidate's Signature : Invigilator's Signature :
Facsimile Signature Stamp of
Centre Superintendent :

E2



AU=9400. 14 13. 7. Which one of the following compounds shows the presence of intramolecular hydrogen bond? (3) HCN \not{K} h-C= CH₃- $\dot{C}-OH$ (4) Cellulose H_{1} O(1) Concentrated acetic acid 121 H2O2 8. The molar conductivity of a $0.5 \text{ mol}/\text{dm}^3$ of AgNO3 with electrolytic solution conductivity of $5.76 \times 10^{-3} \text{ S cm}^{-1}$ at 298 K is (1) 28.8 S cm²/mol $100 \times K = \lambda_{M}$ (2) 2.88 S cm²/mol $M = \lambda_{M}$ (3) 11.52 S cm²/mol $1000 \times 5.76 \times 10^{5}$ (4) 0.086 S cm²/mol 11.5(4) 0.086 S cm^2 /mol 9.) The decomposition of phosphine (PH3) on tungsten at low pressure is a first-order reaction. It is because the (1) rate of decomposition is very slow (2) rate is proportional to the surface coverage (3) rate is inversely proportional to the surface coverage (4) rate is independent of the surface coverage 10. The coagulation values in millimoles per litre of the electrolytes used for the coagulation of As2S3 are given below : 2 I. (NaCl) = 52, II. $(BaCl_2) = 0.69$, III. $(MgSO_4) = 0.22$ $(. V \propto 1)$ The correct order of their coagulating power is (1) III > I > II (2) I > II > III (3) II > I > III . (4) III > II > I 11. During the electrolysis of molten sodium chloride, the time required to produce 0.10 mol of chlorine gas using a current of JMD/E2 0^{1-2} 0^{1-2} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0^{1} 0

12. How many electrons can fit in the orbital for which n = 3 and l = 1? 12 2 3p. III (1) 14 (3) 6 13. For a sample of perfect gas when its pressure is changed isothermally from p_i to p_f , the entropy change is given by $\overset{\text{(1)}}{=} \Delta S = RT \ln \left(\frac{p_i}{p_f} \right) \qquad \qquad \overset{\text{(1)}}{=} \frac{V_{2^*}}{V_1} \quad \overset{\text{(1)}}{=} \frac{P_i}{P_2}$ (2) $\Delta S = nR \ln \left(\frac{p_f}{p_i}\right) \qquad \Delta S = \frac{g_{rev}}{T}$ (3) $\Delta S = nR \ln \left(\frac{p_i}{p_f}\right) \qquad = -\prod k \frac{f}{T} \ln \frac{v_{X_i}}{T}$ (4) $\Delta S = nRT \ln \left(\frac{p_f}{p_i}\right) \approx = RT \ln \frac{v_{Y_i}}{T}$ 14. The van't Hoff factor (i) for a dilute aqueous solution of the strong electrolyte barium hydroxide is Ba (042, 13 (2) 0 (4) 2 (3) 1 15. The percentage of pyridine (C5H5N) that forms pyridinium ion ($C_5H_5N^+H$) in a 0.10 M aqueous pyridine solution (K_b for 18 0.013% (4) 0.77% = 15" × 1.3. 16. In calcium fluoride, having the fluorite structure, the coordination numbers for calcium ion (Ca2+) and fluoride ion (F) are - 11 4 and 8 (2) 4 and 2 (3) 6 and 6 (4) 8 and 4 17. If the E_{cell} for a given reaction has a negative value, which of the following gives the correct relationships for the values of ΔG° and K_{eq} ? (1) $\Delta G^{\circ} < 0$; $K_{eq} < 1$ K $E_{e} - \Delta G^{\circ} = - \pi K \ln k_c$ (2) $\Delta G^{\circ} > 0$; $K_{eq} < 1$ $\int G^{\circ} > 0; K_{eq} < 1$ \sim (3) $\Delta G^{\circ} > 0; K_{eq} > 1$ (4) $\Delta G^{\circ} < 0; K_{eq} > 1$

1.69


0.1 mol. 0.2 °P.T.O.

1e-> 1mole->9/40.

24. Zinc can be coated on iron to produce 18. Which one of the following is incorrect for galvanized iron but the reverse is not ideal solution? possible. It is because (1) zinc has higher negative electrode potential than iron (2) zinc is lighter than iron (3) $\Delta U_{\text{mix}} = 0$ (3) zinc has lower melting point than iron (4) $\Delta P = P_{obs} - P_{calculated}$ by Raoult's law = 0 (#) zinc has lower negative electrode 19. The solubility of AgCl (s) with solubility potential than iron product 1.6×10^{-10} in 0.1 M NaCl solution 25. The suspension of slaked lime in water is Ag CP = NJ" + CI" would be known as VAT 1.26×10-5 M NG(-> Nat + (1) (1) aqueous solution of slaked lime (2) limewater balout the (3) $1.6 \times 10^{-9} M$ (4) $1.6 \times 10^{-11} M$ $S^{2} = \sqrt{1.6 \times 10^{-10} \times 0}$ $S^{2} = \sqrt{1.6 \times 10^{-10} \times 0}$ XIO (a0+MD (3) quicklime (4) $1.6 \times 10^{-11} M$ = $\sqrt{16 \times 10^{-10} \times 0^{-1}}$ = $\sqrt{16 \times 10^{-10} \times 0^{-1}}$ = $4 \times 10^{-6} \times \sqrt{0^{-1}}$ 20. Suppose the elements X and Y. combine to form two compounds XY₂ and X₃Y₂. When 0.1 mole of XY₂ weighs 10 g and 0.05 mole of X₃Y₂ weighs 9 g, the atomic weights of X and Y are (4) milk of lime 26. The hybridizations of atomic orbitals of nitrogen in NO_2^+ , NO_3^- and NH_4^+ respectively il il HAM are (1) sp^2 , sp and sp^{3} X+27=100 (2) 40, 30 3×27 = 180 X and Y are (2) sp, sp³ and sp² χ (1) 30, 20 $\chi^{[3]}$ sp², sp³ and sp χ sp, sp^2 and sp^3 21. The number of electrons delivered at the cathode during electrolysis by a current of 27. Which of the following fluoro-compounds is 1 ampere in 60 seconds is (charge on electron = 1.60×10^{-19} C) most likely to behave as a Lewis base? (1) 7.48×10^{23} (2) 6×10^{23} (2) 6×10^{23} (2) 6×10^{19} (2) BF3 (1) SiF_A (8) PF2 (4) CF4 (3) 6×10^{20} (4) 3.75×10^{20} 28. Which of the following pairs of ions is isoelectronic and isostructural? $(H) ClO_3^2, SO_3^2 (2) CO_3^{2-}, NO_3^2$ 22. Boric acid is an acid because its molecule (1) combines with proton from water (3) $C10_3^-$, $C0_3^{2-}$ (4) $S0_3^{2-}$, $N0_3^{-}$ molecule (2) contains replaceable H^+ ion K - B - MIn context with beryllium, which one of the (3) gives up a proton 29) following statements is incorrect? (4) accepts OH⁻ from water releasing proton (1) Its hydride is electron-deficient and 23. AlF₃ is soluble in HF only in presence of KF. polymeric. (2) It is rendered passive by nitric acid., It is due to the formation of (1) $K[AIF_3H]$ (2) $K_3[AIF_3H_3]$ (3) It forms Be₂C. (4) Its salts rarely hydrolyze. (3) $K_3[AlF_6]$ (4) AlH_3

JMD/E2

Beth




JMD/E2

5

[P.T.O.

1 mole - 9648.



46. A foreign DNA and plasmid cut by the same	51 The primary producers of the deep-sea hydrothermal vent ecosystem are
restriction endonuclease can be joined to form a recombinant plasmid using	
(1) ligase	(1) coral reefs K
(2) $Eco RI$	(2) green algae 🕊
	(3) chemosynthetic bacteria
(3) Taq polymerase	(4) blue-green algae√
(4) polymerase III	ŝ
47. Which of the following is not a component of downstream processing?	(52.) Which of the following is correct for r -selected species?
Expression	(1) Small number of progeny with large size
	(2) Large number of progeny with small size
(2) Separation	(3) Large number of progeny with large size
(3) Purification	(4) Small number of progeny with small size
(4) Preservation	
48. Which of the following restriction enzymes produces blunt ends?	53. If '+' sign is assigned to beneficial interaction, '-' sign to detrimental and '0' sign to neutral interaction, then the population interaction
(1) Hind III	represented by '+' '-' refers to
(2) Sal I	(1) parasitism + - parasit
(3) Eco RV +	(2) mutualism fagasif
(4) Xho I	(3) amensalism
	(4) commensalism
49. Which kind of therapy was given in 1990 to a four-year-old girl with adenosine deaminase (ADA) deficiency?	54. Which of the following is correctly matched?
(1) Radiation therapy	(1) Stratification—Population
(2) Gene therapy	(2) Aerenchyma—Opuntia
(3) Chemotherapy	(3) Age pyramid—Biome
(4) Immunotherapy	(4) Parthenium hysterophorus—Threat to biodiversity
50. How many hot spots of biodiversity in the world have been identified till date by Norman Myers?	55. Red List contains data or information on (1) marine vertebrates only
43	
(2) 17	(2) all economically important plants
(3) 25	 (3) plants whose products are in international trade
(4) 34	(4) threatened species
(1) 01	
JMD /E2	7 [P.T.O.

1 more > 4648.

62. The term 'polyadelphous' is related to 56. Which one of the following is wrong for (1) calyx fungi? and unicellular (2) gynoecium both are (1) They multicellular. 131 androecium (2) They are eukaryotic. (3) All fungi possess a purely cellulosic cell (4) corolla many plants among Indigofera, wall. How Sesbania, Salvia, Allorm, Albe, mustard; (4) They are heterotrophic. groundnut, radish, gram and furnip have 57. Methanogens belong to stamens with different lengths in their (1) Slime moulds abaceae flowers? Monoadelphous (2) Eubacteria (1) Six (3) Archaebacteria (2) Three (4) Dinoflagellates (3) Four 58. Select the wrong statement. (1) Diatoms are microscopic and float (4) Five passively in water. 64. Radial symmetry is found in the flowers of easily walls of diatoms are The 12) Acti (1) Cassia K destructible. (3) 'Diatomaceous earth' is formed by the Brassica Cale cell walls of diatoms. (4) Diatoms are chief producers in the (3) Trifolium 🗙 (4) Pisum oceans. 59. The label of a herbarium sheet does not 65. Free-central placentation is found in carry information on (1) Citrus (A) height of the plant (2) date of collection 2 Dianthus (3) name of collector (3) Argemone (4) local names -(4) Brassica 60. Conifers are adapted to tolerate extreme 66. Cortex is the region found between environmental conditions because of (1) endodermis and vascular bundle (1) presence of vessels (2) broad hardy leaves (2) epidermis and stele (3) superficial stomata (3) pericycle and endodermis (A) thick cuticle (4) endodermis and pith 61. Which one of the following statements is 67. The balloon-shaped structures called tyloses wrong ? (1) Laminaria and Sargassum are used as (1) are linked to the ascent of sap through food. xylem vessels (2) Algae increase the level of dissolved (2) originate in the lumen of vessels oxygen in the immediate environment.-(3) Algin is obtained from red algae, and (3) characterize the sapwood carrageenan from brown algae. (A) are extensions of xylem parenchyma (4) Agar-agar is obtained from Gelidium and cells into vessels Gracilaria.

Cou

JMD/E2

C. Myz

 69. Select the mismatch. (1) Methanogens—Prokaryotes* (2) Gas vacuoles—Green bacteria (3) Alkaline (4) Low refractive index \$\$ (5) Large central vacuoles—Animal cells* (4) Protists—Eukaryotes* (7) Select the wrong statement. (1) Mycoplasma is a wall-less imicroorganism. (2) Bacterial cell wall is made up of petidoglycan. (3) Pili and fimbriae are mainly involved in motility of bacterial cells. (4) Cyanobacteria lack flagellated cells. (7) A cell organelle containing hydrolytic enzymes is (1) mesosome (2) Incrosome (3) microsome (4) ribosome (3) G₁ phase (3) G₁ phase (3) G₁ phase (3) G₁ phase (4) G₂ phase (5) Glucose-6-phosphate^X. (6) Fructose 1, 6-bisphosphate ^X. (7) Mich of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (4) Pyruvic acid (3) Fructose 1, 6-bisphosphate ^X. (4) Pyruvic acid (3) Microscopic server that leads to the death of fishes. (2) Offspring produced by the asexual reproductive structures are called coospores. (3) Fructose 1, 6-bisphosphate ^X. (4) Pyruvic acid (5) Pyruvic acid (5) Pyruvic acid (6) Pyruvic acid 	 68. A non-proteinaceous enzyme is (1) deoxyribonuclease (2) lysozyme (3) ribozyme (4) ligase 	 74. A few drops of sap were collected by cutting across a plant stem by a suitable method. The sap was tested chemically. Which one of the following test results indicates that it is phloem sap? (1) Absence of sugar X (2) Acidic
 (1) Methanogens—Prokalyotes' (2) Gas vacuoles—Green bacteria (3) Protists—Eukaryotes (4) Protists—Eukaryotes (5) Select the wrong statement. (1) Mycoplasma is a wall-less microorganism. (2) Bacterial cell wall is made up of peptidoglycan. (3) Pili and Gimbriar are mainly involved in motility of bacterial cells. (4) Cyanobacteria lack flagellated cells. (7) A cell organelle containing hydrolytic enzymes is (1) mesosome (2) Ibacome (3) Microsome (4) ribosome (5) S phase (3) G₁ phase (4) G₂ phase (5) S phase (3) G₁ phase (4) G₂ phase (5) S phase (6) Glucose-6-phosphate^K. (7) Fuctose 1,6-bisphosphate ^K. (7) Structose 1,6-bisphosphate^K. (7) Structose 1,6-bisph	69. Select the mismatch.	(3) Alkaline
 (2) Gas vacuoles—Green bacteria (3) Large central vacuoles—Animal cells (4) Protists—Eukaryotes (5) Select the wrong statement. (1) Mycoplasma is a wall-less microorganism. (2) Bacterial cell wall is made up of peptidoglycan. (3) Pili and fimbriae are mainly involved in motility of bacterial cells. (4) Cyanobacteria lack flagellated cells. (4) Cyanobacteria lack flagellated cells. (4) Cyanobacteria lack flagellated cells. (5) Kar and give protection (6) The process which makes major difference between C₃ and C₄ plants is (7) Shase (3) G₁ phase (4) G₂ phase (5) G₁ phase (6) G₂ phase (7) Acetyl CoA (2) Glucose-6-phosphate^K. (3) Fructose 1,6-bisphosphate K (4) Pyruvic acid 	(1) Methanogens—Prokaryotes	(4) Low refractive index ⋠
 (d) Large central vacuoles—Animal cells (e) Protists—Eukaryotes 70. Select the wrong statement. (1) Mycoplasma is a wall-less microorganism. (2) Bacterial cell wall is made up of peptidoglycan. (3) Pili and fimbriae are mainly involved in motility of bacterial cells. (4) Cyanobacteria lack flagellated cells. 71. A cell organelle containing hydrolytic enzymes is (1) mesosome (2) Jacoma (2) Zn- (3) microsome (4) ribosome (5) Sphase (3) G₁ phase (4) G₂ phase 73. Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (4) Acetyl CoA (2) Glucose-6-phosphate^K. (3) Fructose 1,6-bisphosphate K (4) Pyruvic acid 	(2) Gas vacuoles—Green bacteria	75. You are given a tissue with its potential for
 70. Select the wrong statement. (1) Mycoplasma is a wall-less microorganism. (2) Bacterial cell wall is made up of peptidoglycan. (3) Pili and fimbriae are mainly involved in motility of bacterial cells. (4) Cyanobacteria lack flagellated cells. 71. A cell organelle containing hydrolytic enzymes is (1) mesosome (2) IaA and gibberellin (3) Auxin and cytokinin (4) Auxin and abscisic acid (5) Phytochrome is a (6) Cyanobacteria lack flagellated cells. 71. A cell organelle containing hydrolytic enzymes is (1) mesosome (2) IaVa and gibberellin (3) Giportein (4) Lipoprotein (5) Revention (2) Zn. (6) Free (4) Ca (7) Which is essential for the growth of root tip? (1) Mnt. (2) Zn. (3) Fe (4) Ca (3) Fe (4) Ca (4) A can discusse major difference between C3 and C4 plants is (5) S phase (6) G1 phase (7) Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (1) Acetyl CoA (2) Glucose-6-phosphate^N (3) Fructose 1,6-bisphosphate ^N (4) Pyruvic acid 	(3) Large central vacuoles—Animal cells -	of the following pairs of hormones would you add to the medium to secure shoots as well
 1) Mycoplasma is a wall-less microorganism. 2) Bacterial cell wall is made up of peptidoglycan. 3) Pili and fimbriae are mainly involved in motility of bacterial cells. 4) Cyanobacteria lack flagellated cells. 4) Cyanobacteria lack flagellated cells. 71. A cell organelle containing hydrolytic enzymes is (1) mesosome (2) IAA and gibberellin (3) Auxin and cytokinin (4) Auxin and abscisic acid 76. Phytochrome is a (1) chromoprotein (2) flavoprotein (3) Fe (4) Ca (77) Which is essential for the growth of root tip? (1) Mnv- (2) Zn. (3) Fe (4) Ca (78) The process which makes major difference between C₃ and C₄ plants is (4) Fe process which makes major difference between C₃ and C₄ plants is (5) Fe (4) Ca (78) The process which makes major difference between C₃ and C₄ plants is (4) photorespiration (5) Calvin cycle X (4) photorespiration (5) Calvin cycle X (6) G₂ phase (7) Neich of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (4) Acetyl CoA (2) Glucose-6-phosphate X (3) Fructose 1,6-bisphosphate X (4) Pyruvic acid 	70 Select the mrong statement	
 (a) microorganism. (b) Bacterial cell wall is made up of petidoglycan. (c) Bacterial cell wall is made up of petidoglycan. (d) Cyanobacteria lack flagellated cells. (e) Cyanobacteria lack flagellated cells. (f) Suppose (f) Imesosome (g) Sphase (g) G₁ phase (g) G₂ phase (g) G₁ phase (g) G₁ phase (g) G₁ phase (g) G₁ phase (g) G₂ phase (g) Gucose-6-phosphate (f) (g) Fructose 1,6-bisphosphate (f) (h) Pyruvic acid 		
 (2) Bacterial cell wall is made up of peptidoglycan. (3) Pili and fimbriae are mainly involved in motility of bacterial cells. (4) Cyanobacteria lack flagellated cells. (4) Cyanobacteria lack flagellated cells. (4) Cyanobacteria lack flagellated cells. (5) Flytochrome is a (1) chromoprotein (2) flavoprotein (3) glycoprotein (4) lipoprotein (7) Which is essential for the growth of root tip? (1) mesosome (3) microsome (4) ribosome (3) microsome (4) ribosome (3) microsome (4) ribosome (3) microsome (4) ribosome (5) Free (4) Ca (7) Which is essential for the growth of root tip? (1) Mnv (2) Zn. (3) Fe (4) Ca (7) Which is essential for the growth of froot tip? (1) Mnv (2) Zn. (3) Fe (4) Ca (7) Which is essential for the growth of root tip? (1) Mnv (2) Zn. (3) Fe (4) Ca (7) Which is essential for the growth of froot tip? (1) Mnv (2) Zn. (3) Fe (4) Ca (7) Which is essential for the growth of root tip? (1) Mnv (2) Zn. (3) Fe (4) Ca (7) Which is essential for the growth of root tip? (1) Mnv (2) glycolysis (3) Calvin cycle X (4) photorespiration N (7) Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (4) Acetyl CoA (5) Fructose 1,6-bisphosphate X (4) Pyruvic acid (5) Fructose 1,6-bisphosphate X (6) Pyruvic acid 	(-)	
 76. Phytochrome is a (1) chromoprotein (2) flavoprotein (3) priti and fimbriae are mainly involved in motility of bacterial cells. (4) Cyanobacteria lack flagellated cells. 71. A cell organelle containing hydrolytic enzymes is (1) mesosome (2) flavoprotein (3) glycoprotein (4) lipoprotein 77 Which is essential for the growth of root tip? (1) Mntx. (2) Zn. (3) Fe (4) Ca 78 The process which makes major difference between C₃ and C₄ plants is (3) Fe (4) G₂ phase (3) G₁ phase (4) G₂ phase 78 Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (4) Acetyl CoA (2) Glucose-6-phosphate X (3) Fructose 1,6-bisphosphate X (4) Pyruvic acid 		
 (3) Pili and fimbriae are mainly involved in motility of bacterial cells. (4) Cyanobacteria lack flagellated cells. (7) A cell organelle containing hydrolytic enzymes is (1) chromoprotein (2) flavoprotein (3) glycoprotein (4) lipoprotein (7) Which is essential for the growth of root tip? (1) Mnx- (2) Zn- (3) Fe (4) Ca (7) Which is essential for the growth of root tip? (1) Mnx- (2) Zn- (3) Fe (4) Ca (7) Which is essential for the growth of root tip? (1) Mnx- (2) Zn- (3) Fe (4) Ca (7) Which is essential for the growth of root tip? (1) Mnx- (2) Zn- (3) Fe (4) Ca (7) Which is essential for the growth of root tip? (1) Mnx- (2) Zn- (3) Fe (4) Ca (7) Which is essential for the growth of root tip? (1) Mnx- (2) Zn- (3) Fe (4) Ca (7) Which is essential for the growth of root tip? (1) Mnx- (2) Zn- (3) Fe (4) Ca (7) Which is essential for the growth of root tip? (1) Mnx- (2) Zn- (3) Fe (4) Ca (7) Which or the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (1) Water hyacinth, growing in the standing water, drains oxygen from water that leads to the death of fishes. (2) Glucose-6-phosphate^K (3) Fructose 1,6-bisphosphate^K (4) Pyruvic acid 		
 motility of bacterial cells. (4) Cyanobacteria lack flagellated cells. 71. A cell organelle containing hydrolytic enzymes is (1) mesosome (2) flavoprotein (3) glycoprotein (4) lipoprotein (77) Which is essential for the growth of root tip? (1) Mnx. (2) Zn. (3) Fe (4) Ca (3) Fe (4) Ca (3) Fe (4) Ca (4) ribosome (5) Fe (4) Ca (78) The process which makes major difference between C₃ and C₄ plants is (1) M phase (2) glycolysist (C) (3) Calvin cycle (C) (4) photorespiration (5) S phase (3) G₁ phase (4) G₂ phase (5) S phase (6) G₂ phase (78) Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (78) Micho se the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (77) Which one of the following statements is not correct? (1) Water hyacinth, growing in the standing-water, drains oxygen from water that leads to the death of fishes. (2) Offspring produced by the asexual reproductive structures are called cone. (3) Microscopic, motile asexual reproductive structures are called zoospores. (4) Pyruvic acid ' 	(3) Pili and fimbriae are mainly involved in	
 (4) Cyanobacteria lack flagellated cells. (7) A cell organelle containing hydrolytic enzymes is (1) mesosome (2) Iysosome (3) microsome (4) ribosome (7) Which is essential for the growth of root tip? (1) Mnx. (2) Zn. (3) Fe (4) Ca (3) Fe (4) Ca (3) Fe (4) Ca (3) Fe (4) Ca (4) ribosome (5) Fe (4) Ca (6) Fe (4) Ca (7) Which is essential for the growth of front tip? (1) Mnx. (2) Zn. (3) Fe (4) Ca (3) Fe (4) Ca (4) protocess which makes major difference between C₃ and C₄ plants is (2) glycolysist (2) Glycolysist (2) Glycolysist (3) Calvin cycle (4) photorespiration (5) (3) Calvin cycle (4) photorespiration (7) (4) photorespiration (7) (5) Phase (6) G₂ phase (7) Which one of the following statements is not correct? (1) Water hyacinth, growing in the standing-water, drains oxygen from water that leads to the death of fishes. (2) Offspring produced by the asexual reproductive structures are called clone. (3) Microscopic, motile asexual reproductive structures are called zoospores. (4) Pyruvic acid ' 	motility of bacterial cells.	
 77 Which is essential for the growth of root tip? (1) mesosome (2) Iyosoome (3) microsome (4) ribosome (7) Which is essential for the growth of root tip? (1) Mnx. (2) Zn. (3) Fe (4) Ca (3) Fe (4) Ca (3) Fe (4) Ca (3) Fe (4) Ca (4) G₂ phase (5) S phase (6) G₁ phase (7) Which is essential for the growth of root tip? (1) Mnx. (2) Zn. (3) Fe (4) Ca (3) Fe (4) Ca (4) G₂ phase (5) G₁ phase (6) G₂ phase (7) Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (7) Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (7) Which one of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (7) Which is essential for the growth of root tip? (1) Mnx. (2) Zn. (3) Fe (4) Ca (4) phase (5) Chase (6) Chase (7) Which is essential for the growth of root tip? (1) Mnx. (2) Zn. (3) Fe (4) Ca (4) photorespiration (5) Chase (6) Chase (7) Which one of the following statements is not correct? (1) Water hyacinth, growing in the standing water, drains oxygen from water that leads to the death of fishes. (2) Offspring produced by the asexual reproductive structures are called zoospores. (3) In potato, banana and ginger, the plantlets arise from the internodes present in the modified stem. 	(4) Cyanobacteria lack flagellated cells.	
 enzymes is (1) mesosome (2) lysosome (3) microsome (4) ribosome (7) During cell growth, DNA synthesis takes place in (1) M phase (7) S phase (3) G₁ phase (4) G₂ phase (7) Which is essential for the growth of root tip? (1) M m⁴. (2) Zn. (3) Fe (4) Ca (3) Fe (4) Ca (3) Calvin cycle X (4) gp hase (5) G₁ phase (6) G₂ phase (7) Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (1) Acetyl CoA (2) Glucose-6-phosphate X (3) Fructose 1,6-bisphosphate X (4) Pyruvic acid (5) Fructose 1,6-bisphosphate X (4) Pyruvic acid (5) Fructose 1,6-bisphosphate X (6) Pyruvic acid (7) Which is essential for the growth of root tip? (1) Mm⁴. (2) Zn. (3) Fe (4) Ca (3) Fe (4) Ca (3) Calvin cycle X (4) photorespiration (2) glycolysis (3) Calvin cycle X (4) photorespiration (4) Pyruvic acid (5) Offspring produced by the asexual reproductive structures are called zoospores. (6) In potato, banana and ginger, the plantlets arise from the internodes present in the modified stem. 	71. A cell organelle containing hydrolytic	(4) lipoprotein
 (1) mesosome (2) lysosome (3) microsome (4) ribosome (72. During cell growth, DNA synthesis takes place in (1) M phase (1) M phase (1) M phase (2) glycolysisch (3) Calvin cyclex (4) G₂ phase (73. Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (4) Acetyl CoA (2) Glucose-6-phosphatex (3) Fructose 1,6-bisphosphate X (4) Pyruvic acid 		(77) Which is essential for the growth of root tip?
 (3) microsome (4) ribosome (78) The process which makes major difference between C₃ and C₄ plants is (78) The process which makes major difference between C₃ and C₄ plants is (78) The process which makes major difference between C₃ and C₄ plants is (78) The process which makes major difference between C₃ and C₄ plants is (78) The process which makes major difference between C₃ and C₄ plants is (78) The process which makes major difference between C₃ and C₄ plants is (78) The process which makes major difference between C₃ and C₄ plants is (79) Which orespiration (2) glycolysist (3) Calvin cycle (4) photorespiration (79) Which one of the following statements is not correct? (1) Water hyacinth, growing in the standingwater, drains oxygen from water that leads to the death of fishes. (2) Offspring produced by the asexual reproductive structures are called clone. (3) Kicroscopic, motile asexual reproductive structures are called zoospores. (4) Pyruvic acid (5) Fructose 1,6-bisphosphate (4) Pyruvic acid 	(1) mesosome	
 (4) ribosome (4) ribosome (4) ribosome (72. During cell growth, DNA synthesis takes place in (1) M phase (2) glycolysis (3) G₁ phase (4) G₂ phase (73. Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (7) Acetyl CoA (2) Glucose-6-phosphate (3) Fructose 1,6-bisphosphate (4) Pyruvic acid 	2 lysosome	(3) Fe (4) Ca
 (4) ribosome (4) ribosome (4) ribosome (72. During cell growth, DNA synthesis takes place in (1) M phase (2) glycolysis (3) G₁ phase (4) G₂ phase (73. Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (7) Acetyl CoA (2) Glucose-6-phosphate (3) Fructose 1,6-bisphosphate (4) Pyruvic acid 	(3) microsome	The process which makes major difference
 72. During cell growth, DNA synthesis takes place in M phase M phase S phase G₁ phase G₂ phase 73. Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? M Acetyl CoA Glucose-6-phosphate Fructose 1,6-bisphosphate Fructose 1,6-bisphosphate 74. The production of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? M Acetyl CoA Glucose-6-phosphate Fructose 1,6-bisphosphate Pyruvic acid 		
 place in (1) M phase (2) S phase (3) G₁ phase (4) G₂ phase 73. Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (1) Water hyacinth, growing in the standing water, drains oxygen from water that leads to the death of fishes. (2) Offspring produced by the asexual reproduction are called clone. (3) Fructose 1,6-bisphosphate * (4) Pyruvic acid (5) Fructose 1,6-bisphosphate * (6) Pyruvic acid 		respiration
 (1) M phase (3) Calvin cyclex (4) S phase (4) G₂ phase (73. Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (4) Acetyl CoA (2) Glucose-6-phosphatex (3) Fructose 1,6-bisphosphate x (4) Pyruvic acid (3) Calvin cyclex (4) photorespiration (5) Which one of the following statements is not correct? (1) Water hyacinth, growing in the standing water, drains oxygen from water that leads to the death of fishes. (2) Offspring produced by the asexual reproduction are called clone. (3) Calvin cyclex (4) photorespiration (5) Which one of the following statements is not correct? (1) Water hyacinth, growing in the standing water, drains oxygen from water that leads to the death of fishes. (2) Offspring produced by the asexual reproductive structures are called clone. (3) Fructose 1,6-bisphosphate x (4) Pyruvic acid 		(2) glycolysis
 (4) S phase (3) G₁ phase (4) G₂ phase 73. Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (4) Acetyl CoA (2) Glucose-6-phosphate (3) Fructose 1,6-bisphosphate (4) Pyruvic acid (5) Fructose 1,6-bisphosphate (6) Photohesphate (7) Which one of the following statements is not correct? (1) Water hyacinth, growing in the standing water, drains oxygen from water that leads to the death of fishes. (2) Offspring produced by the asexual reproduction are called clone. (3) Fructose 1,6-bisphosphate (4) Pyruvic acid (5) Photohesphate (7) Which one of the following statements is not correct? (1) Water hyacinth, growing in the standing water, drains oxygen from water that leads to the death of fishes. (2) Offspring produced by the asexual reproductive structures are called clone. (3) Fructose 1,6-bisphosphate (4) Pyruvic acid (5) Photohesphate (6) Photohesphate (7) Which one of the following statements is not correct? (1) Water hyacinth, growing in the standing water, drains oxygen from water that leads to the death of fishes. (2) Offspring produced by the asexual reproductive structures are called clone. (3) Microscopic, motile asexual reproductive structures are called zoospores. (4) Pyruvic acid 		
 (3) G₁ phase (4) G₂ phase 73. Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (1) Water hyacinth, growing in the standing water, drains oxygen from water that leads to the death of fishes. (2) Offspring produced by the asexual reproduction are called clone. (3) Fructose 1,6-bisphosphate × (4) Pyruvic acid ' (5) Fructose 1,6-bisphosphate × (6) Pyruvic acid ' 		(4) photorespiration
 (4) G₂ phase (73. Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (1) Water hyacinth, growing in the standing water, drains oxygen from water that leads to the death of fishes. (2) Offspring produced by the asexual reproduction are called clone. (3) Fructose 1,6-bisphosphate * (4) Pyruvic acid * 	V. ·	
 73. Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (1) Acetyl CoA (2) Glucose-6-phosphate (3) Fructose 1,6-bisphosphate (4) Pyruvic acid 		
 73. Which of the following biomolecules is common to respiration-mediated breakdown of fats, carbohydrates and proteins? (1) Acetyl CoA (2) Glucose-6-phosphate (*) (3) Fructose 1,6-bisphosphate (*) (4) Pyruvic acid ' 	(4) G ₂ phase	
 of fats, carbohydrates and proteins? (1) Acetyl CoA (2) Glucose-6-phosphate (3) Fructose 1,6-bisphosphate (4) Pyruvic acid (5) Fructose 1,6-bisphosphate (6) Pyruvic acid (7) In potato, banana and ginger, the plantlets arise from the internodes present in the modified stem. 	73. Which of the following biomolecules is	leads to the death of fishes.
 (1) Acetyl CoA (2) Glucose-6-phosphate (3) Fructose 1,6-bisphosphate (4) Pyruvic acid (5) Microscopic, motile asexual reproductive structures are called zoospores. (4) Pyruvic acid (5) Microscopic, motile asexual reproductive structures are called zoospores. (4) In potato, banana and ginger, the plantlets arise from the internodes present in the modified stem. 	common to respiration-mediated breakdown	
 (2) Glucose-6-phosphate (3) Fructose 1,6-bisphosphate (4) Pyruvic acid (5) Fructose 1,6-bisphosphate (6) Pyruvic acid (7) In potato, banana and ginger, the plantlets arise from the internodes present in the modified stem. 	/	
 (2) Glucose-b-phosphate (3) Fructose 1,6-bisphosphate (4) Pyruvic acid (5) Fructose 1,6-bisphosphate (6) Pyruvic acid (7) In potato, banana and ginger, the plantlets arise from the internodes present in the modified stem. 		
 (3) Fructose 1,6-bisphosphate (4) Pyruvic acid (5) Pyruvic acid (6) Pyruvic acid (7) Pyruvic acid (8) Pyruvic acid (9) Pyruvic acid (10) Pyruvic		
		plantlets arise from the internodes
JMD /E2 9 [P.T.O.	(4) Pyruvic acid	present in the modified stem.
	JMD/E2	9 [P.T.O.

JMD/E2

1 mole - 9648.

80. Which one of the following generates new	85. Taylor conducted the experiments to prove
genetic combinations leading to variation?	semiconservative mode of chromosome
(1) Nucellar polyembryony	replication on
(2) Vegetative reproduction	(1) E. coli
(3) Parthenogenesis	(2) Vinca rosea
(4) Sexual reproduction	VIST Vicia faba
81. Match Column-I with Column-II and	(4) Drosophila melanogaster
select the correct option using the codes given below :	86. The mechanism that causes a gene to move from one linkage group to another is called
Column—I Column—II	(1) crossing-over K
a. Pistils fused (i) Gametogenesis	(2) inversion \checkmark
together	(3) duplication
b. Formation of (ii) Pistillate gametes	(#) translocation
c. Hyphae of higher (iii) Syncarpous	V
Ascomycetes	87. The equivalent of a structural gene is
d. Unisexual female (iv) Dikaryotic	(1) recon
flower	(2) muton ×
Codes :	vist cistron
a b c d	(4) operon
(iii) (i) (iv) (ii)	88. A true breeding plant is
(2) (iv) (iii) (i) (ii) (3) (ii) (i) (iv) (iii)	(1) always homozygous recessive in its genetic constitution
(3) (11) (1) (1V) (111) (4) (11) (11) (1V) (111)	(2) one that is able to breed on its own
82. In majority of angiosperms	(3) produced due to cross-pollination among unrelated plants
(1) a small central cell is present in the embryo sac X	(4) near homozygous and produces offspring of its own kind
(2) egg has a filiform apparatus	
(3) there are numerous antipodal cells K	89. Which of the following rRNAs acts as structural RNA as well as ribozyme in
(A) reduction division occurs in the megaspore mother cells	bacteria?
83. Pollination in water hyacinth and water lily is	(1) 5.8 S rRNA
83. Pollination in water hyacinth and water highs brought about by the agency of	(2) 5 S rRNA χ'
(1) bats	(3) 18 S rRNA
12) water	(4) 23 S TRNA
(3) insects or wind	90. Stirred-tank bioreactors have been designed
(4) birds	for
84. The ovule of an angiosperm is technically equivalent to	(1) ensuring anaerobic conditions in the culture vessel
(1) megaspore	(2) purification of product
127 megasporangium	(3) addition of preservatives to the product
(3) megasporophyll	(A) availability of oxygen throughout the
(4) megaspore mother cell	process
a se heards)

91.	A molecule that can act as a genetic material must fulfill the traits given below, except	96. Match Column—I with Column—II and select the correct option using the codes
	(1) it should provide the scope for slow changes that are required for evolution	Column—I Column—II
	(2) it should be able to express itself in the form of 'Mendelian characters'	a. Citric acid (i) Trichoderma b. Cyclosporin A (ii) Clostridium
	(3) it should be able to generate its replicat	c. Statins (iii) Aspergillus d. Butyric acid (iv) Monascus
^	it should be unstable structurally and chemically	Codes: a b c d
92.	DNA-dependent RNA polymerase catalyzes transcription on one strand of the DNA which is called the	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
	(1) antistrand	(4) (i) (iv) (ii) (iii)
	(2) template strand	(97) Biochemical Oxygen Demand (BOD) may not
	(3) coding strand	be a good index for pollution for water bodies receiving effluents from
	(4) alpha strand	(1) sugar industry → Molasser → Hig (2) domestic sewagek
~~		(2) domestic sewagek
93.	Interspecific hybridization is the mating of	(3) dairy industry — 0
	(1) more closely related individuals within same breed for 4-6 generations	(4) petroleum industry98. The principle of competitive exclusion was
	(2) animals within same breed without having common ancestors	stated by (1) Verhulst and Pearl
,	(3) two different related species	(2) C. Darwin
	(4) superior males and females of different	(4) MacArthur
-	breeds	(99) Which of the following National Parks is
94.	Which of the following is correct regarding AIDS causative agent HIV?	(1) Dachigam National Park, Jammu &
anti	(1) HIV does not escape but attacks the <u>acquired immune response</u> . T- ymphod	Kashmir (2) Keibul Lamjao National Park, Manipur (3) Bandhavgarh National Park, Madhya
	(2) HIV is enveloped virus containing one molecule of single-stranded RNA and one molecule of reverse transcriptase.	 (3) Bandhavgarh National Park, Madhya Pradesh (4) Eaglenest Wildlife Sanctuary, Arunachal Pradesh
	(3) HIV is enveloped virus that contains two identical molecules of single-stranded RNA and two molecules of reverse transcriptase.	 100. A lake which is rich in organic waste may result in (1) mortality of fish due to lack of oxygen (2) increased population of aquatic
-	(4) HIV is unenveloped retrovirus.	organisms due to minerals χ (3) drying of the lake due to algal bloom χ
(95)	Among the following edible fishes, which one is a marine fish having rich source of omega-3 fatty acids?	(4) increased population of fish due to lots of nutrients α
	(1) Mackerel	101. The highest DDT concentration in aquatic food chain shall occur in
	(2) Mystus	(1) cel
		(2) phytoplankton
	(3) Mangur	seagull .
	(4) Mrigala	(4) crab

1 mole - >9648.



102. Which of the following sets of diseases is 108. Oxidative phosphorylation is caused by bacteria?

- (1) Herpes and influenzak
- (2) Cholera and tetanus
- (3) Typhoid and smallpox V
- (4) Tetanus and mumps K
- 103. Match Column-I with Column-II for housefly classification and select the correct option using the codes given below :
 - Column-I a. Family

Column-II

- (i) Diptera (ii) Arthropoda b. Order c. Class
 - (iii) Muscidae
- d. Phylum . (iv) Insecta
- Codes :
- d h C a (iiii) (iv) (ii) (i) (1)(ii) -(2) (iii) (i) (iv) (3) (iii) (ii) (iv) (i)
- (4) (iv) (iii) (ii) (i)
- 104. Choose the correct statement.
 - (1) All Pisces have gills covered by an operculum. K
 - (2) All mammals are viviparous.
 - (3) All cyclostomes do not possess jaws and paired fins.
 - (4) All reptiles have a three-chambered heart. X
- 105. Study the four statements (A-D) given below and select the two correct ones out of them :
 - A. Definition of biological species was given by Ernst Mayr.v
 - B. Photoperiod does not affect reproduction in plants.X
 - C. Binomial nomenclature system was given by R. H. Whittake
 - D. In unicellular organisms, reproduction is synonymous with growth.
 - The two correct statements are
 - (1) A and B (2) B and C
 - (3) C and D A and D

In male cockroaches, sperms are stored in which part of the reproductive system?

- (1) Vas deferens
- (2) Seminal vesicles
- . (3) Mushroom glands

(4) Testes v


107. Smooth muscles are

(1) voluntary, spindle-shaped, uninucleate

- (2) involuntary, fusiform, non-striated
 - (3) voluntary, multinucleate, cylindrical
 - (4) involuntary, cylindrical, striated &

JMD/E2

- - (1) formation of ATP by energy released from electrons removed during substrate oxidation
 - (2) formation of ATP by transfer o phosphate group from a substrate to ADP
 - (3) oxidation of phosphate group in ATP
 - (4) addition of phosphate group to ATP
 - 109. Which of the following is the least likely to be involved in stabilizing the three-dimensiona folding of most proteins?
 - (1) Ester bonds
 - (2) Hydrogen bonds√
 - (3) Electrostatic interaction
 - Hydrophobic interaction
 - 110. Which of the following describes the giver graph correctly?



- (1) Exothermic reaction with energy A in absence of enzyme and B in presence of enzyme
- (2) Endothermic reaction with energy A in presence of enzyme and B in absence of enzyme
- Exothermic reaction with energy A in presence of enzyme and B in absence of enzyme
- Endothermic reaction with energy A in absence of enzyme and B in presence of enzyme
- 111. When cell has stalled DNA replication fork, which checkpoint should be predominantly activated?

(1) Both G_2/M and M 121 G1/S (3) G_2/M (4) M

DNA Replication

112. Match the stages of meiosis in Column-I to | 117. Name the ion responsible for unmasking of active sites for myosin for cross-bridge their characteristic features in Column-II activity during muscle contraction. and select the correct option using the codes given below : (1) Potassium (2) Calcium Column-I Column-II Magnesium (4) Sodium a. Pachytene Pairing of homologous (i) 118. Name the blood cells, whose reduction in chromosomes number can cause clotting disorder, leading b. Metaphase I (ii) Terminalization of to excessive loss of blood from the body. chiasmata (1) Thrombocytesc. Diakinesis (iii) Crossing-over takes place (2) Erythrocytes Chromosomes align at d. Zygotene (iv) (3) Leucocytes equatorial plate -292 (4) Neutrophils Codes : 119. Name a peptide hormone which acts mainly b d c a on hepatocytes, adipocytes and enhances (1) (iv) (iii) (ii) (i) 2 cellular glucose uptake and utilization. n (2) (iii) (iv) (ii) (i) (2) Insulin Y (1) Gastrin (iii) (3)(i) (iv) (ii) Glucagon (4) Secretin (4) (ii) (iv) (iiii) (i) 120. Osteoporosis, an age-related disease of 113. Which hormones do stimulate the skeletal system, may occur due to and production of pancreatic juice (1) accumulation of uric acid leading to bicarbonate? inflammation of joints (1) Insulin and glucagon (2) immune disorder affecting neuro-(2) Angiotensin and epinephrine muscular junction leading to fatigue (3) Gastrin and insulin (3) high concentration of Ca⁺⁺ and Na⁺ (A) Cholecystokinin and secretin (4) decreased level of estrogen 114. The partial pressure of oxygen in the alveoli 121. Serum differs from blood in of the lungs is . (1) lacking antibodies (1) less than that of carbon dioxidek (2) lacking globulins (2) equal to that in the blood . (3) lacking albumins (3) more than that in the blood (4) lacking clotting factors (4) less than that in the blood 122. Lungs do not collapse between breaths and Choose the correct statement. some air always remains in the lungs which (1) Receptors do not produce graded can never be expelled because potentials. (1) pressure in the lungs is higher than the (2) Nociceptors respond to changes in atmospheric pressurer pressure. (2) there is a negative pressure in the lungsk (3) Meissner's corpuscles are thermo-(3) there is a negative intrapleural pressure receptors. pulling at the lung walls .(4) Photoreceptors in the human eye are (4) there is a positive intrapleural pressure λ depolarized during darkness and become hyperpolarized in response to the light 123. The posterior pituitary gland is not a 'true' stimulus. endocrine gland because 116. Graves' disease is caused due to (1) it secretes enzymes (A) hypersecretion of adrenal gland X (2) it is provided with a duct (2) hyposecretion of thyroid gland , 137 it only stores and releases hormones (3) hypersecretion of thyroid gland & (4) it is under the regulation of hypo-(4) hyposecretion of adrenal gland thalamus JMD/E2 [P.T.O. 13

1 mole - 79648.

(124) The part of nephron involved in active	130. Several hormones like hCG, hPL, estroger
reabsorption of sodium is	progesterone are produced by
(1) descending limb of Henle's loop	(1) pituitary
(2) distal convoluted tubule DCT	(2) ovary
(3) proximal convoluted tubule	
(4) Bowman's capsule v	vo placenta
125. Which of the following is hormone- releasing IUD?	(4) fallopian tube
(1) Cu7	131. If a colour-blind man marries a woman while home-blind for normal colour vision the
(2) LNG-20	is homozygous for normal colour vision, the probability of their son being colour-blind is
(3) Multiload 375	
(4) Lippes loop	$(1) 1 \qquad (2) 0 \qquad X \qquad$
126. Which of the following is incorrect regarding vasectomy?	(3) 0.5 (4) 0.75 XXC (XY)
(1) Irreversible sterility Keversibility	132. Genetic drift operates in
(2) No sperm occurs in seminal liuidvi	(1) slow reproductive population
No sperm occurs in epididymis &	(2) small isolated population
(4) Vasa deferentia is cut and tied 127. Embryo with more than 16 blastomeres	
formed due to in vitro fertilization is	
transferred into	.,
(1) cervix	133. In Hardy-Weinberg equation, the frequence
12) uterus	of heterozygous individual is represented by
(3) fallopian tube	(1) q^2 (2) p^2
(4) fimbriae	(3) 2pq (4) pq
128. Which of the following depicts the correct	
pathway of transport of sperms? (1) Efferent ductules \rightarrow Rete testis \rightarrow Vas	(134) The chronological order of human evolutic
deferens \rightarrow Epididymis	from early to the recent is
(2) Rete testis \rightarrow Efferent ductules \rightarrow	(XI) Australopithecus → Homo habilis
\checkmark Epididymis \rightarrow Vas deferens	Ramapithecus >> Homo erectus
(3) Rete testis \rightarrow Epididymis \rightarrow Efferent	(2) Australopithecus \rightarrow Ramapithecus
ductules \rightarrow Vas deferens	Homo habilis -> Homo erectus
(4) Rete testis → Vas deferens → Efferent ductules → Epididymis	(3) Ramapithecus \rightarrow Australopithecus
129. Match Column—I with Column—II and	Homo habilis -> Homo erectus
select the correct option using the codes	
given below :	Australopithecus \rightarrow Homo erectus
Column—I Column—II	135. Which of the following is the correct
a. Mons pubis(i) Embryo formation	sequence of events in the origin of life?
b. Antrum (ii) Sperm	I. Formation of protobionts
c. Trophectoderm (iii) Female external	
∧ genitalia	II. Synthesis of organic monomers*
d. Nebenkern / (iv) Graafian follicle	III. Synthesis of organic polymers /
Codes :	IV. Formation of DNA-based genetic system
a , b c d	(1) II, III, IV, I
(1) (i) (iv) (iii) (ii)	(2) I, II, III, IV
(2) (iii) · (iv) (ii) · (i)	(3) I, III, II, IV
	(4) II, III, I, IV
(4) (iii) (i) (iv) (ii)	
JMD/E2	14

*

1=1f=so21+1=1 400 s.f=so21+1=1 $\begin{array}{c} 45 \\ +1 \\ +1 \\ 400 \\ 5 \\ \end{array} \begin{array}{c} f \\ f \\ f \\ 400 \\ \end{array} \begin{array}{c} f \\ f \\ f \\ f \\ 400 \\ \end{array} \begin{array}{c} f \\ f \\ f \\ f \\ 400 \\ \end{array} \begin{array}{c} f \\ f \\ f \\ f \\ 400 \\ \end{array} \begin{array}{c} f \\ f \\ f \\ f \\ 400 \\ \end{array} \begin{array}{c} f \\ f \\ f \\ f \\ 400 \\ \end{array}$ O 2 A person can see clearly objects only when 140. If an electron in a hydrogen atom jumps they lie between 50 cm and 400 cm from his from the 3rd orbit to the 2nd orbit, it emits eyes. In order to increase the maximum a photon of wavelength λ . When it jumps distance of distinct vision to infinity, the type from the 4th orbit to the 3rd orbit, the and power of the correcting lens, the person corresponding wavelength of the photon has to use, will be 1 10-2 will be $\frac{\lambda'}{\lambda} = \frac{5}{26} \frac{\chi_{14}}{\frac{16}{16}} + \frac{1}{\lambda} = R_{14} \left[\frac{1}{4} - \frac{1}{9} \right]$ $\frac{\lambda'}{\lambda} = \frac{5}{26} \frac{\chi_{14}}{\frac{16}{16}} + \frac{1}{\lambda} = R_{14} \left[\frac{1}{4} - \frac{1}{9} \right]$ (1) convex, +0.15 diopter 24(1) $\frac{20}{13}\lambda$ (2) convex, +2.25 diopter $\frac{150}{700}$ (8) concave, -0.25 diopter (3) $\frac{9}{16}\lambda$ (4) concave, -0.2 diopter 50 141. The half-life of a radioactive substance is A linear aperture whose width is 0.02 cm is 30 minutes. The time (in minutes) taken placed immediately in front of a lens of focal between 40% decay and 85% decay of the length 60 cm. The aperture is illuminated ubstance is (2) 15 In $\frac{60^{4}}{15} = -\lambda(\Delta T)$ (4) 45 2 109 = $\frac{102}{30} \Delta T$ same radioactive substance is normally by a parallel beam of wavelength 5×10^{-5} cm. The distance of the first dark 60 band of the diffraction pattern from the (3) 30 centre of the screen is 5×105× 6000 AD 0102 00 - 84 0502 - 3× 5×10 15001 - 3× 5×10 200942. (n) 0.15 cm For CE transistor amplifier, the audio signal voltage across the collector resistance of 2 kΩ (2) 0.10 cm is 4 V. If the current amplification factor of 100 = 3(3) 0.25 cm the transistor is 100 and the base resistance is $1 k\Omega$, then the input signal voltage is (4) 0.20 cm P FB= iEti (1) 15 mV (2) 10 mV BLOXIS 138. Electrons of mass m with de-Broglie (3) 20 mV (4) 30 mV wavelength λ fall on the target in an X-ray tube. The cutoff wavelength (λ_0) of the 143. The given circuit has two ideal diodes emitted X-ray is connected as shown in the figure below. The $\frac{f^2}{2m} = \frac{b^2}{2m/2} = \frac{Kc}{4}$ (1) $\lambda_0 = \lambda$ current flowing through the resistance R_1 will be Na $\sqrt{2} \lambda_0 = \frac{2mc\lambda^2}{c}$ 0.027-(3) $\lambda_0 = \frac{2h}{mc}$ (4) $\lambda_0 = \frac{2m^2c^2\lambda^3}{h^2}$ V $u = \frac{1}{f}$ $P = \frac{1}{100}$ cmi 2Ω R_1 10 V Z2Q5V D 139. Photons with energy 5 eV are incident 2.5 A (1) 3·13 A on a cathode C in a photoelectric cell. The maximum energy of emitted photoelectrons (3) 10.0 A (4) 1.43 A is 2 eV. When photons of energy 6 eV are incident on C, no photoelectrons will reach 144. What is the output Y in the following circuit, the anode A, if the stopping potential of A when all the three inputs A, B, C are first relative to C is 0 and then 1? (1) -3 V B. C+ (3) +4 V (1) 1, 1 (2) 0, 1 (3) 0, 0 (4) 1, 0 JMD/E2 15 [P.T.O.

1 mole - 9648.

$$2 = \frac{1}{2} \times \frac{10}{1000} \times (1600 - \sqrt{2} - 100) = \frac{1000}{100} =$$

(1)
$$\sqrt{\frac{Gc}{h^{3/2}}}$$
 $\sqrt{2} \frac{\sqrt{hG}}{c^{3/2}} ML^{2}T^{-1}M^{3}L^{5}T^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L^{-3}L$

146. Two cars P and Q start from a point at the same time in a straight line and their positions are represented by $x_P(t) = at + bt^2$ and $x_0(t) = ft - t^2$. At what time do the cars have the same velocity? have the same velocity: (1) $\frac{f-a}{2(1+b)}$ (2) $\frac{a-f}{1+b}$ VQ= f-2t (3) $\frac{a+f}{2(b-1)}$ (4) $\frac{a+f}{2(1+b)}$ f-2t= 0+2bt f-0=(2b+2)

47) In the given figure, $a = 15 \text{ m/s}^2$ represents the total acceleration of a particle moving in the clockwise direction in a circle of radius R = 2.5 m at a given instant of time. The speed of the particle is,

$$\begin{array}{c} 2 - 5 \\ V = \frac{15}{92} \\ 3 \\ 2 - 5 \\ 2 - 5 \\ 2 - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ - 5 \\ -$$

148. A rigid ball of mass m strikes a rigid wall at V². 60° and gets reflected without loss of speed as shown in the figure below. The value of impulse imparted by the wall on the ball TS. Will be

 $\frac{1200 = \sqrt{2}}{10 \sqrt{12} = \sqrt{2} \frac{4 \cos n s^{-1}}{\sqrt{10}}}$ A bullet of mass 10 g moving horizonts

with a velocity of 400 m s⁻¹ strikes a wood block of mass 2 kg which is suspended b light inextensible string of length 5 m. A result, the centre of gravity of the block found to rise a vertical distance of 10 c The speed of the bullet after it emerges (horizontally from the block will be

(1)
$$160 \text{ m s}^{-1} = \frac{1}{2} \times 10^{2}$$

(2) $100 \text{ m s}^{-1} = \frac{1}{2} \times 10^{2}$
(3) $80 \text{ m s}^{-1} = \frac{1}{200} \times 10^{2}$
(4) 120 m s^{-1}
 $V = \sqrt{200}$
 $V = \sqrt{200}$

 $V_p = \alpha + 245_{150}$. Two identical balls A and B having velociti of 0.5 m/s and -0.3 m/s respectively colli elastically in one dimension. The velocities B and A after the collision respectively will 1 (1) 0.3 m/s and 0.5 m/s

- (2) -0.5 m/s and 0.3 m/s (3) 0.5 m/s and -0.3 m/s (x -0.3 m/s and 0.5 m/s
- **151.** A particle moves from a point $(-2\hat{i} + 5\hat{j})$ $(4\hat{j}+3\hat{k})$ when a force of $(4\hat{i}+3\hat{j})$ N applied. How much work has been done 1 the force? ~ A

(1) 2 J
$$21 - 1$$

(1) 2 J $41 - 5$
(2) 8 J $8 - 3$
(3) 11 J $40 - 1 \times 10 \times 10$
(4) 5 J $20 - 3$
(3) $12 + 5 - 1 \times 10 \times 10$
(600 - $\sqrt{2}$
(7) $21 - 1 \times 10 \times 10$
(600 - $\sqrt{2}$
(7) $21 - 1 \times 10 \times 10$
(600 - $\sqrt{2}$
(7) $21 - 1 \times 10 \times 10$
(7) $12 \times 10 \times 10$
(7) $12 \times 10 \times 10^{-1}$
(8) $11 = 1 \times 10^{-1} \times 10^{-1}$
(9) $5 = 1 \times 10^{-1} \times 10^{-1} \times 10^{-1}$
(9) $5 = 1 \times 10^{-1} \times 10^{-1} \times 10^{-1}$
(9) $5 = 1 \times 10^{-1} \times 10^{-1} \times 10^{-1}$
(9) $5 = 1 \times 10^{-1} \times 10^{-1} \times 10^{-1}$
(9) $5 = 1 \times 10^{-1} \times 10^{-1} \times 10^{-1}$
(9) $5 = 1 \times 10^{-1} \times 10^{-1} \times 10^{-1} \times 10^{-1}$
(9) $5 = 1 \times 10^{-1} \times 10^{-1} \times 10^{-1} \times 10^{-1}$
(9) $5 = 1 \times 10^{-1} \times 10^{-1}$

152. Two rotating bodies A and B of masses with moments of inertia I_A an A) have equal kinetic energy (If L_A and L_B be their angula respectively, then

(1)
$$L_A > L_B$$
 $L^2 \times T$ $L_B > L_A$
(2) $L_A = \frac{L_B}{2}$ Ψ

don y n

$$(3) \quad L_A = 2L_B$$


153. A solid sphere of mass m and radius R is 15 rotating about its diameter. A solid cylinder of the same mass and same radius is also rotating about its geometrical axis with an angular speed twice that of the sphere. The ratio of their kinetic energies of rotation

(2) 2:3 137 1:5

- (4) 1:4
- 154. A light rod of length l has two masses m_1 and m_2 attached to its two ends. The moment of inertia of the system about an axis perpendicular to the rod and passing through the centre of mass is

$$(\mathcal{M}_{1}) \frac{m_{1}m_{2}}{m_{1}m_{2}} l^{2} \qquad (\mathcal{M}_{1}) \frac{m_{1}m_{2}}{m_{1}+m_{2}} l^{2} \qquad (\mathcal{M}_{1}) \frac{m_{1}m_{2}}{m_{1}+m_{2}} l^{2} \qquad (\mathcal{M}_{1}) \frac{m_{1}m_{2}}{m_{1}m_{2}} l^{2} \qquad (\mathcal{M}_{2}) l^{2} \qquad (\mathcal{M}_{2}) l^{2} \qquad (\mathcal{M}_{2}) l^{2} \qquad (\mathcal{M}$$

155. Starting from the centre of the earth having radius R, the variation of g (acceleration due to gravity) is shown by



156. A satellite of mass m is orbiting the earth (of radius R) at a height h from its surface. The total energy of the satellite in terms of g_0 , the value of acceleration due to gravity at the earth's surface, is

(1)
$$-\frac{2mg_0R^2}{R+h}$$
 $-\frac{6nMm}{ng_0R^2}$ $q(R+h)$
(2) $\frac{mg_0R^2}{2(R+h)}$ $q(R+h)$ $q(R+h)$
 $\sqrt{(3)} -\frac{mg_0R^2}{2(R+h)}$ $T = k \left[\frac{5T}{2} - \frac{3T}{2} \right]$
(4) $\frac{2mg_0R^2}{R+h}$ $T = k \left[\frac{3T}{2} - \frac{3T}{2} \right]$
 $K = 1$
 $K = 3T_2$
 $K = 3T_2$

157. A rectangular film of liquid is extended
from
$$(4 \text{ cm} \times 2 \text{ cm})$$
 to $(5 \text{ cm} \times 4 \text{ cm})$. If the
work done is 3×10^{-4} J, the value of the
surface tension of the liquid is
(1) $8 \cdot 0 \text{ Nm}^{-1}$ (2) $0 \cdot 250 \text{ Nm}^{-1}$
(3) $0 \cdot 125 \text{ Nm}^{-1}$ (4) $0 \cdot 2 \text{ Nm}^{-1}$ (5)
158. Three liquids of densities ρ_1, ρ_2 and ρ_3 (with
 $\rho_1 > \rho_2 > \rho_3$), having the same value of
surface tension *T*, rise to the same height
in three identical capillaries. The angles of
contact θ_1, θ_2 and θ_3 obey
(1) $\pi > \theta_1 > \theta_2 > \theta_3 > \frac{\pi}{2}$
(2) $\frac{\pi}{2} > \theta_1 > \theta_2 > \theta_3 > \frac{\pi}{2}$
 $(3) 0 \le \theta_1 < \theta_2 < \theta_3 < \frac{\pi}{2}$
 $(3) 0 \le \theta_1 < \theta_2 < \theta_3 < \frac{\pi}{2}$
 $(4) (3) 0 \le \theta_1 < \theta_2 < \theta_3 < \frac{\pi}{2}$
 $(4) (4) (\frac{\pi}{4} < \theta_1 < \theta_2 < \theta_3 < \theta_3) < \frac{\pi}{2}$
 $(4) (4) (\frac{\pi}{4} < \theta_1 < \theta_2 < \theta_3 < \pi$

159. Two identical bodies are made of a material for which the heat capacity increases with temperature. One of these is at 100 °C, while the other one is at 0 °C. If the two bodies are brought into contact, then, assuming no heat loss, the final common temperature is $(-1)^{(1)} 0 °C$ (1) 0 °C(2) 50 °C(2) 50 °C(3) 7-100 = 50 °C

- more than 50 °C
- (4) less than 50 °C but greater than 0 °C
- 160. A body cools from a temperature 3T to 2T in 10 minutes. The room temperature is T. Assume that Newton's law of cooling is applicable. The temperature of the body at the end of next 10 minutes will be


$$\begin{array}{c} (1) & T \\ (1) & T \\ (3) & T \\ (3) & \frac{3}{2}T \\ (3$$

61. One mole of an ideal monatomic gas undergoes a process described by the equation PV^3 = constant. The heat capacity of the gas during this process is

(1)
$$R$$
 (2) $\frac{3}{2}R$
(3) $\frac{5}{2}R$ (4) $2R$

$$C = Cv + \frac{R}{N-1} \quad [P.T.O.]$$
$$= \frac{3R}{2} - \frac{R}{2} - R$$

1 more >4648



A parallel-plate capacitor of area A, plat separation d and capacitance C is filled wit four dielectric materials having dielectric constants k_1 , k_2 , k_3 and k_4 as shown in th figure below. If a single dielectric material is to be used to have the same capacitance C is this capacitor, then its dielectric constant is given by

$$(1) \quad \frac{1}{k} = \frac{1}{k_1} + \frac{1}{k_2} + \frac{1}{k_3} + \frac{3}{2k_4} = \frac{C'}{3} + \frac{2}{k_1 + k_2} + \frac{1}{k_3} + \frac{3}{2k_4} = \frac{C'}{3} + \frac{2}{k_1 + k_2} + \frac{1}{k_3} + \frac{3}{2k_4} + \frac{1}{2k_4} +$$

169. The potential difference $(V_A - V_B)$ betwee the points A and B in the given figure is


$$V_{A} \xrightarrow{2\Omega} \frac{3V}{+} 1\Omega = \frac{1}{2} \frac{V_{E}}{A} \xrightarrow{I = 2} A \xrightarrow{I = 2} \frac{1}{4} \frac{V_{E}}{+} \xrightarrow{I = 1} \frac{V_{E}}{-} \xrightarrow{V_{A} - \frac{1}{4} - 3} \xrightarrow{V_{E}} \xrightarrow{V_{A} - \frac{1}{4} - 3} \xrightarrow{V_{E}} \xrightarrow{V_{A} - \frac{1}{4} - 3} \xrightarrow{I = 2} \xrightarrow{I =$$


170. A filament bulb (500 W, 100 V) is to be use in a 230 V main supply. When a resistance is connected in series, it works perfectly an the bulb consumes 500 W. The value of R is

(1) 13
$$\Omega$$
 , $\gamma q = (2) 230 \Omega$, 20Ω
(3) 46 $\Omega_{Q} q = (4) 26 \Omega$ $q = \sqrt{20} q$

171. A long wire carrying a steady current is ben into a circular loop of one turn. The magnet field at the centre of the loop is B. It is the bent into a circular coil of n turns. The magnetic field at the centre of this coil n turns will be 2 TUr = 0.5

(1) $2n^2B$ (2) nB $B = \mu_{\bar{\iota}}\bar{\iota}$ (2) n^2B (4) 2nB $2p\chi$ B'= n3





Test Booklet Code

JMD

This Test Booklet contains 20 pages.

7538948


ZZ

Do not open this Test Booklet until you are asked to do so.

Important Instructions :

- The Answer Sheet is inside this Test Booklet. When you are directed to open the Test Booklet, take out the Answer Sheet and fill in the particulars on Side-1 and Side-2 carefully with blue/black ballpoint pen only.
- The test is of 3 hours duration and Test Booklet contains 180 questions. Each question carries 4 marks. For each correct response, the candidate will get 4 marks. For each incorrect response, one mark will be deducted from the total score. The maximum marks are 720.
- 3. Use Blue/Black Ballpoint Pen only for writing particulars on this page/marking responses.
- 4. Rough work is to be done on the space provided for this purpose in the Test Booklet only.
- 5. On completion of the test, the candidate must hand over the Answer Sheet to the Invigilator before leaving the Room/Hall. The candidates are allowed to take away Test Booklet only with them.
- 6. The CODE for this Test Booklet is **ZZ**. Make sure that the CODE printed on **Side-2** of the Answer Sheet is the same as that on this Test Booklet. In case of discrepancy, the candidate should immediately report the matter to the Invigilator for replacement of both the Test Booklet and the Answer Sheet.
- 7. The candidate should ensure that the Answer Sheet is not folded. Do not make any stray marks on the Answer Sheet. Do not write your Roll No. anywhere else except in the specified space in the Test Booklet/Answer Sheet.
- 8. Use of white fluid for correction is not permissible on the Answer Sheet.
- 9. Each candidate must show on demand his/her Admit Card to the Invigilator.
- 10. No candidate, without special permission of the Superintendent or Invigilator, would leave his/her seat.
- 11. The candidates should not leave the Examination Hall without handing over their Answer Sheet to the Invigilator on duty and sign the Attendance Sheet twice. Cases where a candidate has not signed the Attendance Sheet second time will be deemed not to have handed over the Answer Sheet and dealt with as an unfair means case.
- 12. Use of Electronic/Manual Calculator is prohibited.
- 13. The candidates are governed by all Rules and Regulations of the Board with regard to their conduct in the Examination Hall. All cases of unfair means will be dealt with as per Rules and Regulations of the Board.
- 14. No part of the Test Booklet and Answer Sheet shall be detached under any circumstances.
- 15. The candidates will write the correct Test Booklet Code as given in the Test Booklet/Answer Sheet in the Attendance Sheet.

Name of the Candidate (in Capitals) : ABHILASHA RHATT
Roll Number (in Figures) : 3720068
(in Words) : Eight come seventy two taken sinty eight.
Centre of Examination (in Capitalis): KENDRIVA VIDYALAYA, MATHIRATRIALA, PENEADUN
Centre of Examination (in Capitals) : KENDRIVA VIDYALAYA, HATHIRAR KALA, PEREADUN Candidate's Signature : Different A. Antherent Signature :
Facsimile Signature Stamp of Centre Superintendent :
E4



1. A (solid) sphere of mass *m* and radius *R* is
rotating about its diameter. A solid cylinder
of the same mass and same radius is also
rotating about its geometrical axis with an
angular speed twice that of the sphere. The
ratio of the kinetic energies of rotation
(Esphere / Ecylinder) will be
41 1:5
(2) 1:4
(4) 2:3
A light rod of length *l* has two masses *m*₁ and
*m*₂ attached to its two ends. The moment
of inertia of the system about an axis
perpendicular to the rod and passing
through the centre of mass is
$$J = M_2$$

(1) $\frac{m_1 + m_2}{m_1 m_2} l^2$ (2) $(m_1 + m_2) l^2$
(3) $\sqrt{m_1 m_2} l^2$ (4) $\frac{m_1 m_2}{m_1 + m_2} l^2$
3. Starting from the centre of the earth having
radius *R*, the variation of *g* (acceleration due
to gravity) is shown by
(1) $\frac{1}{0} \int \frac{1}{R - r} e^{\frac{1}{R}} \int \frac{1}{R - r} e^{\frac{1}{R}} e^{\frac{1}{R}$

R

T

J

 $w_{2} = \frac{1}{2}mR^{2} \qquad kRal / lef(2)$ $w_{2} = \frac{1}{2}mR^{2} (\frac{1}{2}mR^{2})(2w)$

- A rectangular film of liquid is extended from $(4 \text{ cm} \times 2 \text{ cm})$ to $(5 \text{ cm} \times 4 \text{ cm})$. If the work done is 3×10^{-4} J, the value of the surface tension of the liquid is
 - (1) 0.125 Nm^{-1} (2) 0.2 Nm^{-1}
 - (3) 8.0 Nm^{-1} (4) 0.250 Nm^{-1}
- Three liquids of densities ρ_1 , ρ_2 and ρ_3 (with $\rho_1 > \rho_2 > \rho_3$), having the same value of surface tension T, rise to the same height in three identical capillaries. The angles of contact θ_1 , θ_2 and θ_3 obey $h \ge \tau uso$ (1) $0 \le \theta_1 < \theta_2 < \theta_3 < \frac{\pi}{2}$ (29)

$$2 + (2) \frac{\pi}{2} < \theta_1 < \theta_2 < \theta_3 < \pi \quad (50) = (10)^{-2}$$

$$(3) \pi > \theta_1 > \theta_2 > \theta_3 > \frac{\pi}{2} \quad (50) < (2)^{-2}$$

$$(4) \frac{\pi}{2} > \theta_1 > \theta_2 > \theta_3 \ge 0 \quad (50)^{-2} \quad (50)^{-2}$$

$$(\theta | < 0)^{-2} < 0^{-2}$$

- Two identical bodies are made of a materia for which the heat capacity increases with temperature. One of these is at 100 °C, while the other one is at 0 °C. If the two bodies are brought into contact, then, assuming no heat loss, the final common temperature is
 - (1) more than 50 °C
 - (2) less than 50 °C but greater than 0 °C
 - (3) 0°C
 - (4) 50 °C
- A body cools from a temperature 3T to 2 in 10 minutes. The room temperature is 7 Assume that Newton's law of cooling i applicable. The temperature of the body a the end of next 10 minutes will be

$$\begin{array}{c} (1) \quad \frac{3}{2} T_{pv} N = cons^{+}(2) \quad \frac{4}{3} T \quad C = Cv + \frac{R}{1 - N} \\ (3) \quad \frac{T}{2} \quad \frac{3R}{2} + \frac{R}{(-2)} \quad (4) \quad \frac{7}{4} T \quad C = \frac{3}{2} R + \frac{R}{1 - N} \\ (-2) \quad (-2) \quad$$

One mole of an ideal monatomic ga undergoes a process described by th equation PV^3 = constant. The heat capacit of the gas during this process is

$$(1) \frac{5}{2}R \frac{3R}{2} - \frac{1}{2}(2) 2R \quad \mathcal{G} = \frac{4}{R} \frac{1}{R}$$

$$(1) \frac{5}{2}R \frac{3R}{2} - \frac{1}{2}(2) 2R \quad \mathcal{G} = \frac{4}{R}$$

$$(1) \frac{5}{2}R \frac{3R}{2} - \frac{1}{2}(2) 2R \quad \mathcal{G} = \frac{4}{R}$$

$$(1) \frac{3}{2}R \frac{3R}{2} - \frac{3}{2}R - \frac{1}{2}R = \frac{1}{2}$$

$$= -\frac{2}{2}R - \frac{1}{2}R - \frac{1}{2}R = \frac{1}{2}R - \frac{1}{2}R - \frac{1}{2}R - \frac{1}{2}R = \frac{1}{2}R - \frac{$$

$$\begin{array}{c} y = TPA \\ y = A_{1} - b_{1} = (0 \times 15^{2}) - (0 \times 16^{2}) \\ \hline AA \\ \hline BA = A_{1} - b_{1} = (0 \times 15^{2}) - (0 \times 16^{2}) \\ \hline AA \\ \hline BA = A_{1} - b_{1} = (0 \times 15^{2}) - (0 \times 16^{2}) \\ \hline AA \\ \hline BA = A_{1} - b_{1} = (0 \times 16^{2}) - (0 \times 16^{2}) \\ \hline AA \\ \hline BA = A_{1} - b_{1} = (0 \times 16^{2}) - (0 \times 16^{2}) \\ \hline AA \\ \hline BA = A_{1} - b_{1} = (0 \times 16^{2}) - (0 \times 16^{2}) \\ \hline AA \\ \hline BA = A_{1} - b_{1} = (0 \times 16^{2}) - (0 \times 16^{2}) \\ \hline AA \\ AA \\ \hline A$$

$$w = P + (\cos 0 - \cos 60)$$

 $w = P + (1 - \frac{1}{2}) = \frac{P}{2} = -\frac{MB}{2}$

25833

20. A bar magnet is hung by a thin cotton thread in a uniform horizontal magnetic field and is in equilibrium state. The energy required to rotate it by 60° is W. Now the torque required to keep the magnet in this new MB-2W position is (2) $\frac{\sqrt{3}W}{2}$.

(1)
$$\sqrt{3}W$$

(3) $\frac{2W}{\sqrt{3}}$

21. An electron is moving in a circular path under the influence of a transverse magnetic field of 3.57×10^{-2} T. If the value of e/m is 1.76×10^{11} C/kg, the frequency of revolution of the electron is


(4) $\frac{W}{\sqrt{2}}$

- (2) 62·8 MHz (1) 100 MHz
 - (4) 1 GHz (3) 6.28 MHz
- 22. Which of the following combinations should be selected for better tuning of an L-C-R circuit used for communication?
 - (1) $R = 25 \Omega$, L = 2.5 H, $C = 45 \mu$ F
 - (2) $R = 15 \Omega$, L = 3.5 H, $C = 30 \mu$ F
 - (3) $R = 25 \Omega, L = 1.5 H, C = 45 \mu F$
 - (4) $R = 20 \Omega$, L = 1.5 H, $C = 35 \mu$ F

23. A uniform magnetic field is restricted within a region of radius r. The magnetic field

changes with time at a rate $\frac{dB}{dt}$. Loop 1 of

radius R > r encloses the region r and loop 2 of radius R is outside the region of magnetic field as shown in the figure below. Then the e.m.f. generated is



C= MBSIN 00 = MBG E=MBSin0 24. The potential differences across

resistance, capacitance and inductance are 80 V, 40 V and 100 V respectively in an L-C-R circuit. The power factor of this circuit is

the

= 700 h

(1)
$$0.5$$
 (2) 0.8
(3) 1.0 (4) 0.4

A 100 Ω resistance and a capacitor of 100 Ω 25 reactance are connected in series across a 220 V source. When the capacitor is 50% charged, the peak value of the displacement VI=JR R 13 (2) 4.4 A = 2×10 ×1×12 current is

(1) 11 A (4) 2·2 A (3) 11√2 A = 2×10 V

25. Two identical glass ($\mu_g = 3/2$) equiconvex lenses of focal length f each are kept in contact. The space between the two lenses is filled with water ($\mu_w = 4/3$). The focal length =2×1txu of the combination is (2) 4f/3

(1) f

- (4) f/3(3) 3f/4
- 27. An air bubble in a glass slab with refractive index 1.5 (near normal incidence) is 5 cm deep when viewed from one surface and 3 cm deep when viewed from the opposite face. The thickness (in cm) of the slab is
 - (2) 12 (1) 10
 - (4) 8 · (3) 16
- 28. The interference pattern is obtained with, two coherent light sources of intensity ratio n.

$$\begin{aligned} \varphi = \frac{1}{2} \quad \varphi = \frac{1}{2} \quad \varphi = \frac{1}{2} \quad (\chi + \zeta) + (\chi + \zeta) +$$

•

$$\frac{WL}{W^{-1}L^{2}} = \frac{P/L}{P} \frac{W^{-1}L^{2}}{W^{-1}L^{2}} = \frac{P/L}{P}$$
(38. Planck's constant (h), speed of light in constants (0) and herefore a randomental constants (10) are three and anomatic constants (10) are three and anomatic constants (10) are three and demension of ingh?
(1) $\frac{dRO}{dr^{2}}$ (2) $\sqrt{\frac{Nc}{G}} \cdot \frac{M}{R}$
(3) $\frac{NC}{R}$ (1) $\frac{dRO}{dr^{2}}$ (4) $\frac{dRO}{dr^{2}}$
(1) $\frac{dRO}{dr^{2}}$ (4) $\frac{dRO}{dr^{2}}$
(5) $\frac{1}{2} (\frac{1}{R} - \frac{1}{R})$
(6) $\frac{1}{2} (\frac{1}{R} - \frac{1}{R})$ (2) $\frac{dr}{dr} + \frac{1}{R}$
(3) $\frac{1}{2} (\frac{1}{R} - \frac{1}{R})$ (4) $\frac{dr}{dr} + \frac{1}{R}$
(4) $\frac{1}{2} (\frac{1}{R} - \frac{1}{R})$ (4) $\frac{dr}{dr} + \frac{1}{R}$
(5) $\frac{1}{2} (\frac{1}{R} - \frac{1}{R})$ (4) $\frac{dr}{dr} + \frac{1}{R}$
(6) $\frac{1}{R} - \frac{1}{R} - \frac{1}{R} \times \frac{Nc}{R}$
(7) $\frac{1}{2} (\frac{1}{R} - \frac{1}{R})$ (4) $\frac{dr}{dr} + \frac{1}{R}$
(5) $\frac{1}{R} - \frac{1}{R} - \frac{1}{R} \times \frac{Nc}{R}$
(6) $\frac{1}{R} - \frac{1}{R} - \frac{1}{R} \times \frac{Nc}{R}$
(7) $\frac{1}{R} - \frac{1}{R} - \frac{1}{R} \times \frac{1}{R} - \frac{1}{R} \times \frac{1}{R}$
(1) $\frac{1}{R} - \frac{1}{R} - \frac{1}{R} - \frac{1}{R} \times \frac{1}{R} \times \frac{1}{R} - \frac{1}{R} \times \frac{1}{R} - \frac$

i

.


14

Hine I

.

$$\begin{aligned} & (2+1+1+1+1) + 1 \\ & (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+1) + (3+$$


85. The term 'polyadelphous' is related to 79. Which one of the following is wrong for fungi? (1) androecium (All fungi possess a purely cellulosic cell (2) corolla wall. (3) calyx (2) They are heterotrophic. unicellular and (4) gynoecium both (3) They are "multicellular. V many plants among Indigofera, How (4) They are eukaryotic. Sesbania, Salvia, Allium, Albe, mustard, 80. Methanogens belong to groundnut, radish, gram and turnip have stamens with different lengths in their (1) Archaebacteria flowers? (2) Dinoflagellates (1) Four (3) Slime moulds (4) Eubacteria (2) Five (3) Six 81. Select the wrong statement. (1) 'Diatomaceous earth' is, formed by the (4) Three cell walls of diatoms. 87. Radial symmetry is found in the flowers of (2) Diatoms are chief producers in the oceans. (1) Trifolium (3) Diatoms are microscopic and float (2) Pisum passively in water. The walls of diatoms are easily (3) Cassia destructible. Brassica V The label of a herbarium sheet does not 88. Free-central placentation is found in carry information on (1) name of collector (1) Argemone (2) local names V (2) Brassica (2) height of the plant (3) Citrus (4) date of collection (A) Dianthus Conifers are adapted to tolerate extreme environmental conditions because of 89. Cortex is the region found between (1) superficial stomata (1) pericycle and endodermis (2) thick cuticle (2) endodermis and pitho (3) presence of vessels (3) endodermis and vascular bundle (4) broad hardy leaver (#) epidermis and stele ~ 84. Which one of the following statements is SYBK wrong ? The balloon-shaped structures called tyloses 90 (A) Algin is obtained from red algae, and (1) characterize the sapwood carrageenan from brown algae? (2) Agar-agar is obtained from Gelidium and (2) are extensions of xylem parenchyma cells into vessels Gracilaria. (3) Laminaria and Sargassum are used as (3) are linked to the ascent of sap through food. xylem vessels (4) Algae increase the level of dissolved (4) originate in the lumen of vessels oxygen in the imprediate environment. JMD/E4



12

JMD/E4

 115. A molecule that can act as a genetic material must fulfill the traits given below, scept (1) it should be unstable structurally and chemically (2) it should be unstable structurally and chemically (3) it should be unstable structurally and chemically (4) it should be able to express itself in the form of Mendelian cheracters' (1) coding strand (2) alpha strand (3) antistrand (4) alpha strand (5) superior males and females of different breeds. (2) superior males and females of different breeds. (4) animism within as me breed without having common ancestors. (1) anise within as me breed without having common ancestors. (1) Which of the following is correct regarding ADB causative agent HV? (2) HIV is unerveloped rirus that contains two molecule of singe-stranded RNA and two molecules of reverse transcriptase. (3) Among the following either finds, within as and feaster of the sequence of the seq	115. A molecule that can act as a genetic materia	120 Match Column I with Column
(i) it should be able to expert to its replica (ii) it should provide the scope for slow changes that are required for evolution (ii) it should be able to express itself in the form of Mendelian characters' (ii) Andependent RNA polymerase catalyzes transcription on one strand of the DNA which is called the (i) coding strand (ii) coding strand (iii) coding strand (ii) coding strand (iii) coding strand (iii) coding strand (ii) coding strand (iii) coding strand (iiii) coding strand (iii) coding strand (iii) codin	must fulfill the traits given below, except	
19. It should be (instable) structurally and changes that are required for evolution10.Cyclospoin A11.(3) it should provide the scope for slow changes that are required for evolution form of Mendelian characters'(a)(b)(c)(c)(4) it should be able to express itself in the form of Mendelian characters'(c)(c)(c)(c)(1) coding strand (2) alph a strand(c)(c)(c)(c)(c)(2) antistrand (3) antistrand (4) template strand(c)(c)(c)(c)(c)(1) coding strand (2) superior males and females of different breeder(c)(c)(c)(c)(c)(1) animals within same breed for 4-6 generationas/ (4) animals within same breed without having common ancestors/(c)(c)(c)(c)(c)(1)(c)(c)(c)(c)(c)(c)(c)(c)(2)HIV does not escape but attacks the acquired immune response K(c)(c)(c)(c)(c)(1)Manger and two molecules of reverse transcriptase/(c)(c)(c)(c)(c)(2)HIV does not escape but attacks the acquired immune response/K(c)(c)(c)(c)(c)(1)Manger and two molecules of reverse transcriptase/(c)(c)(c)(c)(c)(3)Mencello of single-stranded RNA and one molecule of single-stranded RNA and one molecule of reverse transcriptase/(c)(c)(c)(c)(1)Manger <b< td=""><td>(1) it should be able to generate its replica</td><td>given below :</td></b<>	(1) it should be able to generate its replica	given below :
10. Consistent of the source of a slow changes that are required for evolution of Mendelian characters. (4) it should be able to express itself in the form of Mendelian characters. (5) BDA-dependent RNA polymerase catalyzes transcription on one strand of the DNA which is called the the analysis of the polymerase catalyzes transcription on one strand of the DNA which is called the the analysis of the polymerase catalyzes transcription on one strand of the DNA dependent RNA polymerase catalyzes transcription on one strand of the DNA dependent RNA and two different related species (1) two different related species (2) superior males and females of different breeds for 4-6 generations. (3) more closely related individuals within same breed without having common ancestor. (4) HIV is enveloped virus that contains two identical molecules of single-stranded RNA and two molecules of reverse transcriptases. (2) HIV does not escape but attacks the acquired immune response. (4) HIV is enveloped virus containing one molecule of reverse transcriptase. (5) HIV does not escape but attacks the acquired immune response. (1) Mangur $f' = \int_{-1}^{1} (f_{-1} f_{-1}) f_{-1} = \int_{-1}^{1} (f_{-1} f_{-1}) f_{-1} = \int_{-1}^{1} (f_{-1} f_{-1}) f_{-1} = f_{-1} f$	(2) it should be unstable structurally and	
(4) it should provide the scope for solution (4) it should be able to express itself in the form of Mendelian characters' (5) DAA-dependent RNA polymerase catalyzes transcription on one strand of the DNA (2) alpha strand (3) antistrand (4) attistrand (3) antistrand (3) antistrand (4) template strand (3) antistrand (3) antistrand (4) template strand (5) superior males and females of different breeds' (2) superior males and females of different breeds' (3) more closely related individuals within same breed for 4-6 generations. (4) animals within same breed without having common ancestors' (4) Minks common ancestors' (5) Which of the following is correct regarding AIDS causative agent HIV? (4) HIV is enveloped virus that contains two identical molecules of single-stranded RNA and two molecules of reverse transcriptase. (2) HIV is enveloped virus that contains two identical meleode without having common ancestorse (4) HIV is enveloped virus that contains two identical meleode without having be following eible fishes, which or eas a marine fish having rich source of molecule of reverse transcriptase? (1) Mangur (2) Mingula (2) Mingula (3) Mackerel (4) Mystus (4) Mystus (5) HIV des (5) HU for enveloped virus containing one molecule of reverse transcriptase? (4) Mingula (5) Among the following eible fishes, which or eas a marine fish having rich source of molecule of reverse transcriptase? (4) Mingula (5) Mingula (5) Mingula (5) Mingula (6) Mingula (6) Mingula (7) Mingula (7) Mangur (7) Mingula (7) M		b. Cyclosporin A (iii) Clostridium
(4) it should be able to express itself in the form of Mendelian characters' (4) it should be able to express itself in the form of Mendelian characters' (5) DNA-dependent RNA polymerase catalyzes transcription on one strand of the DNA which is called the (1) coding strand (3) antistrand (4) template strand (5) antistrand (6) template strand (7) template strand (6) the total strand (7) template strand (7) template strand (8) antistrand (9) template strand (9) template strand (1) dairy industry (2) superior males and females of different breeds (2) superior males and females of different breeds (4) animals within same breed without having common ancestors (4) animals within same breed without having common ancestors (2) HIV is enveloped virus that contains two identical molecules of superstranded RNA and two molecules of reverse transcriptase. (2) HIV is enveloped virus containing one molecule of single-stranded RNA and two molecules of reverse transcriptase. (3) Antong the following edible fishes, which one is a marine fish having rich source of (1) Mangur (2) Mirgala (3) Mirgala (4) Mystus (4) Mystus (5) Mackerel (4) Mystus (5) Mackerel (4) Mystus (5) Mackerel (4) Mystus (5) Mirgala (5) Mirgala (6) Mirgala (6) Mirgala (7)		c. Stating (iii) Aspergillus
form of Mendelian characters' 116. DNA-dependent RNA polymerase catalyzes transcription on one strand of the DNA which is called the (1) coding strand (2) alpha strand (3) antistrand (4) template strand 117. Interspecific hybridization is the mating of (4) two different related species \checkmark (2) superior males and females of different threeds: (3) more closely related individuals within same breed for 4-6 generations. (4) animals within same breed without having common ancestors: (4) Hill is enveloped virus that contains two identical molecules of single-stranded RNA and two molecules of reverse transcriptase. (2) HIV is unerveloped retrovirus. (3) HIV does not escape but attacks the acquired immune response. (4) HIV is enveloped virus containing one molecule of single-stranded RNA and one molecule of single-stranded RNA		(N) Monuscus
 116. DNA-dependent RNA polymerase catalyzes transcription on one strand of the DNA which is called the which is called the iteration of the the DNA (i) consistent of the DNA (ii) (iii) (iv) (ii) (iii) (iv) (iii) (iii) (iv) (iii) (iii) (iv) (ii) (iv) (i	form of 'Mendelian characters'	
12. The replanet is the mating of the DNA which is called the (1) coding strand (2) (i) (ii) (iii) (iii) (3) antistrand (4) template strand (4) template strand (5) antistrand (6) the officient related species (7) three specific hybridization is the mating of M1 two different related species (2) superior males and females of different breeds (3) more closely related individuals within same breed for 4-6 generations. (4) animals within same breed without having common ancestors. (4) animals within same breed without having common ancestors. (5) Which of the following is correct regarding AIDS causative agent HIV? (4) HIV is enveloped virus that contains two identical molecules of single-stranded RNA and two molecules of reverse transcriptase. (2) HIV is enveloped virus containing one molecule of reverse transcriptase. (3) Among the following edible fishes, which on is a marine fish having rich source of omega-3 fatty acids? (4) Magur (5) Magala (6) HIV is guale fish having rich source of omega-3 fatty acids? (1) Mangur (2) Mrigala (3) Migala (4) (4) Mystus (4) Mystus (5) (1 - 2h) = $\mathcal{L}(1 - 2h) = \mathcal{L}(1 - 2h) = $		
which is called the (1) coding strand (2) alpha strand (3) antistrand (4) template strand (3) antistrand (4) template strand (5) template strand (6) template strand (7) template strand (7) template strand (8) more closely related individuals within same breed for 4-6 generations. (9) which of the following is correct regarding AIDS causative agent HIV? (1) HIV is enveloped virus that contains two (1) HIV is enveloped virus that contains two (2) HIV is enveloped virus that contains two (3) HIV does not escape but attacks the acquired immune response. (4) HIV is enveloped virus containing one molecule of reverse transcriptase. (4) HIV is enveloped virus containing one molecule of reverse transcriptase. (5) HIV does not escape but attacks the acquired immune response. (1) Mangur (2) Mirgala (3) Mille Matcerel (4) Mystus (4) Mostus (4) Mostus (5) MID/E4 (5) Contained (5) Alternation of a quatic (4) Mystus (5) MID/E4 (5) Contained (5) Alternation of a quatic (6) Mystus (6) Mystus (7) Contained (7) Alternation of a quatic (7) Mackerel (7) Mackerel (7) Mackerel (7) Mackerel (7) Mackerel (7) Mackerel (7) Margala (7) Among the following chilbe fishes, which ore is a marine fish having rich source of (7) Mackerel (7) Mackerel (7) Mackerel (7) Mackerel (7) Mackerel (7) Mackerel (7) Mackerel (7) Margala (7) Alternation of a quatic (7) Alternation of a quati	transcription on one strand of the DNA	
 (2) alpha strand (3) antistrand (4) template strand (4) template strand (5) antistrand (6) the specific hybridization is the mating of the template strand (7) two different related species / (8) superior males and females of different breeds / (9) superior males and females of different breeds / (10) animals within same breed without having common ancestors / (11) Which of the following is correct regarding AIDS causative agent HIV? (12) HIV is enveloped virus that contains two molecules of single-stranded RNA and two molecules of reverse transcriptase. (13) HIV does not escape but attacks the acquired immune response. X (14) HIV is enveloped virus containing one molecule of reverse transcriptase. (15) HIV is enveloped virus containing one molecule of reverse transcriptase. (16) HIV is enveloped virus containing one molecule of single-stranded RNA and two molecules of single-stranded RNA and one molecule of single-stranded RN		
(a) anisistrand (b) anisistrand (c) a	(1) coding strand	
(i) antistrand (i) template strand (ii) dairy industry (i) dai	(2) alpha strand	Ber Serie (DOD) may (MOV)
(4) template strand (4) template strand (1) dairy industry (2) superior males and females of different breeds (3) more closely related individuals within same breed for 4-6 generations (4) animals within same breed without having common ancestors (4) animals within same breed without having common ancestors (5) Which of the following is correct regarding ADDS causative agent HIV? (4) HIV is enveloped virus that contains two identical molecules of single-stranded RNA and two molecules of reverse transcriptase. (2) HIV is unenveloped retrovirus. (3) HIV does not escape but attacks the acquired immune response. (4) HIV is enveloped virus containing one molecule of ingle-stranded RNA and one molecule of single-stranded RNA and one molecule of reverse transcriptase. (1) Mangur (2) Mrigala (3) Mackerel (4) Mystus (4) Mystus (5) $f = f(1 - 2k)$ (4) Mystus (5) $f = f(1 - 2k)$ (5) $f = f(1 - 2k)$ (4) $f = f(1 - 2k)$ (5) $f = f(1 - 2k)$ (5) $f = f(1 - 2k)$ (6) $f = f(1 - 2k)$ (7) $f = f(1 - 2k)$ (7) $f = f(1 - 2k)$ (8) $f = f(1 - 2k)$ (9) $f = f(1 - 2k)$ (10) $f = f(1 - 2k)$ (11) $f = f(1 - 2k)$ (12) $f = f(1 - 2k)$ (13) $f = f(1 - 2k)$ (14) $f = f(1 - 2k)$ (15) $f = f(1 - 2k)$ (15) $f = f(1 - 2k)$ (16) $f = f(1 - 2k)$ (17) $f = f(1 - 2k)$ (17) $f = f(1 - 2k)$ (18) $f = f(1 - 2k)$ (19) $f = f(1 - 2k)$ (10) $f = f(1 - 2k)$ (11) $f = f(1 - 2k)$ (12) $f = f(1 - 2k)$ (13) $f = f(1 - 2k)$ (14) $f = f(1 - 2k)$ (15) $f = f(1 - 2k)$ (15) $f = f(1 - 2k)$ (16) $f = f(1 - 2k)$ (17) $f = f(1 - 2k)$ (17) $f = f(1 - 2k)$ (18) $f = f(1 - 2k)$ (19) $f = f(1 - 2k)$ (19) $f = f(1 - 2k)$ (10) $f = f(1 - 2k)$ (10) $f = f(1 - 2k)$ (10) $f = f(1 - 2k)$ (11) $f = f(1 - 2k)$ (12) $f = f(1 - 2k)$ (13) $f = f(1 - 2k)$ (14) $f = f(1 - 2k)$ (15) $f = f(1 - 2k)$ (15) $f = f(1 - 2k)$ (16	(3) antistrand	
117. Interspecific hybridization is the mating of (17) two different related species $\sqrt{2}$ (2) superior males and females of different breeds $\sqrt{2}$ (3) more closely related individuals within same breed for 4-6 generations $\sqrt{2}$ (4) animals within same breed without having common ancestors $\sqrt{2}$ (4) animals within same breed without having common ancestors $\sqrt{2}$ (4) animals within same breed without having common ancestors $\sqrt{2}$ (4) Mich of the following is correct regarding AIDS causative agent HIV? (5) Which of the following National Parks is 70 home to the famous musk deer or hangul? (1) Bandhavgarh National Park, Madhya Pradesh National Park, Madhya Pradesh National Park, Madhya Pradesh National Park, Madhya Pradesh National Park, Manipur $\frac{1}{7}$ (2) HIV is unenveloped virus containing one molecule of single-stranded RNA and one molecule of ingerest transcriptase. (4) HIV is enveloped virus containing one molecule of ingele-stranded RNA and one molecule of reverse transcriptase. (1) Mangur (2) Mirgala $\frac{1}{2} = \frac{9}{(\frac{1}{1} \frac{1}{2k_{o}})} = \frac{1}{2}$ (1) Mangur (2) Mirgala $\frac{1}{2} = \frac{9}{(\frac{1}{1} \frac{1}{2k_{o}})} = \frac{1}{2}$ (3) Mirgala $\frac{1}{2} = \frac{9}{(\frac{1}{1} \frac{1}{2k_{o}})} = \frac{1}{2}$ (4) Mixtus $\frac{1}{2} = \frac{9}{(\frac{1}{2} \frac{1}{2k_{o}})} = \frac{1}{2}$ (5) Mirgala $\frac{1}{2} = \frac{9}{(\frac{1}{2} \frac{1}{2k_{o}})} = \frac{1}{2}$ (4) Mixtus $\frac{1}{2} = \frac{9}{(\frac{1}{2} \frac{1}{2k_{o}})} = \frac{1}{2}$ (5) Mixture $\frac{1}{2} = \frac{9}{(\frac{1}{2} \frac{1}{2k_{o}})} = \frac{1}{2}$ (4) phytoplankton $\frac{1}{1} = -\frac{1}{2} \frac{1}{2} \frac{1}{2}$ (5) Relative $\frac{1}{2} = \frac{1}{2} $	(4) template strand	
(4) two different related species $$ (2) superior males and females of different breeds $$ (3) more closely related individuals within same breed for 4-6 generations $$ (4) animals within same breed without having common ancestors $$ (4) animals within same breed without having common ancestors $$ (4) animals within same breed without having common ancestors $$ (4) animals within same breed without having common ancestors $$ (4) animals within same breed without having common ancestors $$ (4) animals within same breed without having common ancestors $$ (4) animals within same breed without having common ancestors $$ (5) Which of the following is correct regarding AIDS causative agent HIV? (4) HIV is enveloped virus that contains two identical molecules of single-stranded RNA and two molecules of reverse transcriptases $$ (2) HIV is unenveloped refforirus. (3) HIV does not escape but attacks the acquired immune response χ (4) HIV is enveloped virus containing one molecule of reverse transcriptases $$ (4) HIV is enveloped virus containing one molecule of reverse transcriptases $$ (5) Mangur $9' = \frac{9}{(+\chi_{2})} = \frac{1}{(+\chi_{2})} = \frac{1}{(+\chi_$	117 Interspecific hybridization is the metion of	(2) petroleum industry
 (2) superior males and females of different breeds (2) (3) more closely related individuals within same breed for 4-6 generations (2) (4) animals within same breed without having common ancestor (2) (4) animals within same breed without having common ancestor (2) (5) Which of the following is correct regarding AIDS causative agent HIV? (4) HIV is enveloped virus that contains two identical molecules of single-stranded RNA and two molecules of reverse transcriptase. (2) HIV is unenveloped retrovirus. (3) HIV does not escape but attacks the acquired immune response. (A) (4) HIV is enveloped virus containing one molecule of single-stranded RNA and one molecule of reverse transcriptase. (1) Mangur (2) (-1) (-1) (-1) (-1) (-1) (-1) (-1) (-1		(c) ougui muusujie
breeds (3) more closely related individuals within same breed for 4-6 generations. (4) animals within same breed without having common ancestor. (5) Which of the following is correct regarding AIDS causative agent HIV? (4) HIV is enveloped virus that contains two identical molecules of single-stranded RNA and two molecules of reverse transcriptase. (2) HIV is unenveloped retrovirus. (3) HIV does not escape but attacks the acquired immune response. (4) HIV is enveloped virus containing one molecule of single-stranded RNA and one molecule of reverse transcriptase? (1) Mangur (2) Mrigala (2) Mingala (3) Mackerel (4) Mystus (4) Mystus (5) MD/E4 (5) Eddewed (6) E. Gause (1) C. Darwin (2) Mink of the following National Park, Madhya Pradesh (4) Keibul Lamjao National Park, Manipur (5) Dachigam National Park, Manipur (4) Keibul Lamjao National Park, Manipur (5) Increased population of fish due to lack of oxygen (1) Mangur (2) Mrigala (4) Mystus (5) Mackerel (4) Mystus (5) MD/E4 (4) Mystus (5) C. Caguenest Widlife Sanctuary, Arunachal Pradesh (6) Dachigam National Park, Manipur (7) Mackerel (6) Increased population of fish due to lack of oxygen (1) Mangur (2) Mrigala (3) MD/E4 (4) Mystus (4) Mystus (5) C. Call the following National Park, Manipur (6) Increased population of fish due to lack of oxygen (1) Seagull (1) Seagull (2) Creab (4) phytoplafikton (5) C. Call the following National Park, Manipur (6) Increased population of aquatic food chain shall occur in (1) seagull (2) Creab (4) phytoplafikton (5) C. C. H. (6) Dachigam National Park, Manipur (7) Mackerel (4) Mystus (6) C. Carwing (7) Mackerel (6) C. Carwing (7) Mackerel (7) Mackerel (8) C. Carwing (8) C. Carwing (9) C. Car		
 (4) animals within same breed without having common ancestors/ (5) Verhulst and Pearl To = 3(27-70) Which of the following is correct regarding AIDS causative agent HIV? (4) HIV is enveloped virus that contains two identical molecules of single-stranded RNA and two molecules of reverse transcriptase. (2) HIV is unenveloped retrovirus. (3) HIV does not escape but attacks the acquired immune response. X (4) HIV is enveloped virus containing one molecule of reverse transcriptase. (4) HIV is enveloped virus containing one molecule of reverse transcriptase. (5) Verhulst and Pearl To = 3(27-70) Which of the following National Parks is having result in To = 17-70 Which of the famous musk deer or hangul? (1) Bandhavgarh National Park, Madhya Pradesh (To+17b = 6 f. 2000 (Control of the second park) (Control of the second park) (Control of the second park. Jammu & 75 (Control of the	breeds	
 (4) animals within same breed without having common ancestors (4) animals within same breed without having common ancestors (5) Vernulst and Pearl To = 3(27-To) To = 3(27-	(3) more closely related individuals within same breed for 4-6 generations.	(1) G. F. Gause (2) MacArthur $\begin{pmatrix} 27-70 \end{pmatrix}$ $=$ $=$ $=$ $=$ $=$ $=$ $=$ $=$ $=$ $=$
Which of the following is correct regarding AIDS causative agent HIV? (1) HIV is enveloped virus that contains two identical molecules of single-stranded RNA and two molecules of reverse transcriptase. (2) HIV is unenveloped retrovirus. (3) HIV does not escape but attacks the acquired immune response. (4) HIV is enveloped virus containing one molecule of single-stranded RNA and one molecule of reverse transcriptase? (4) HIV is enveloped virus containing one molecule of reverse transcriptase? (1) Mangur (1) Mangur (1) Mangur (2) Mrigala (1) Mangur (3) Mackerel (4) Mystus (4) Mystus (5) $\int -2 \int (1 - 2h) \\ (4) Mystus$ (6) $\int -2 \int (1 - 2h) \\ (4) Mystus$ (7) $\int -2 \int (1 - 2h) \\ (4) Mystus$ (8) $\int -2 \int (1 - 2h) \\ (4) Mystus$ (9) $\int -2 \int (1 - 2h) \\ (4) Mystus$ (1) $Margur = \int -2 \int (1 - 2h) \\ (4) Mystus$ (1) $Margur = \int -2 \int (1 - 2h) \\ (4) Mystus$ (2) $\int -2 \int (1 - 2h) \\ (4) Mystus$ (3) $\int -2 \int (1 - 2h) \\ (4) Mystus$ (4) $\int -2 \int (1 - 2h) \\ (4) Mystus$ (5) $\int -2 \int (1 - 2h) \\ (4) Mystus$ (6) $\int -2 \int (1 - 2h) \\ (4) Mystus$ (7) $\int -2 \int (1 - 2h) \\ (4) Mystus$ (7) $\int -2 \int (1 - 2h) \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\ (7) = 0 \\$	(4) animals within same breed without	(3) Verhulst and Pearl $To = 3(27 - To)$ (4) C. Darwin $To = BT - 270$
AIDS causative agent HIV? (1) HIV is enveloped virus that contains two identical molecules of single-stranded RNA and two molecules of reverse transcriptase. (2) HIV is unenveloped retrovirus. (3) HIV does not escape but attacks the acquired immune response. (4) HIV is enveloped virus containing one molecule of single-stranded RNA and one molecule of reverse transcriptase. (4) HIV is enveloped virus containing one molecule of reverse transcriptase. (5) Among the following edible fishes, which one is a marine fish having rich source of omega-3 fatty acids? (1) Mangur $g' = g' + \frac{g}{(+\frac{1}{K_c})} + \frac{1}{(2)}$ (2) Mrigala $g' = g(1 - \frac{2}{K_c}) + \frac{1}{(2)}$ (4) Mystus $g' = g(1 - \frac{2}{K_c}) + \frac{1}{(2)} = \frac{1}{(+\frac{1}{K_c})} + \frac{1}{(2)} = \frac{1}{(+\frac{1}{K_c})} + \frac{1}{(+\frac{1}{K_c})} + \frac{1}{(+\frac{1}{K_c})} = \frac{1}{(+\frac{1}{K_c})} + \frac{1}{(+\frac{1}{K_c})} + \frac{1}{(+\frac{1}{K_c})} + \frac{1}{(+\frac{1}{K_c})} = \frac{1}{(+\frac{1}{K_c})} + \frac{1}{(+1$	6	Which of the following National Parks is
 (1) HIV is enveloped virus that contains two identical molecules of single-stranded RNA and two molecules of reverse transcriptase. (2) HIV is unenveloped retrovirus. (3) HIV does not escape but attacks the acquired immune response. (4) HIV is enveloped virus containing one molecule of single-stranded RNA and one molecule of reverse transcriptase. (4) HIV is enveloped virus containing one molecule of reverse transcriptase. (1) Among the following edible fishes, which one is a marine fish having rich source of omega-3 fatty acids? (1) Mangur (2) Mrigala (3) Mackerel (4) Mystus (4) Mystus (5) Mackerel (4) Mystus (6) MIC/E4 (7) = 9 (1 - 2h) = (R18h) (1) Case (1 - 2h) = (R18h) (1) Case (1 - 2h) = (R18h) (1) Phytoplankton (2) Craba (3) Mackerel (4) Phytoplankton (5) Case (1 - 2h) = (R18h) 		
19 Among the following edible fishes, which one is a marine fish having rich source of omega-3 fatty acids? (1) Mangur (2) Mrigala (3) Mackerel (4) Mystus (4) Mystus (5) $f = 2 \left(1 - \frac{2h}{R^2} \right) = \left(\frac{R_1 R_h}{R_1} \right) = \frac{R_1 R_h}{R_1}$ (6) Dachigam National Park, Jammu & J Xashmir (7) To = 6 f Yamedesh (8) Dachigam National Park, Jammu & J Xashmir (4) Keibul Lamjao National Park, Manipur (4) Keibul Lamjao National Park, Manipur (5) Mathematical may (6) Model and the second park for	(X) HIV is enveloped virus that contains two	101310-01
(1) Mangur $g' = g' (1 - 2h)$ (2) MID/E4 $g' = g' (1 - 2h)$ (3) Markerel $g' = g' (1 - 2h)$ (4) Mystus $g' = g' (1 - 2h)$ (5) Dachigam National Park, Jammu & 3' Kashmir $T_{0} = MT$ (4) Keibul Lamjao National Park, Manipur T_{2} (4) Keibul Lamjao National Park, Manipur T_{2} (5) mortality of the lake due to algal bloom (1) drying of the lake due to algal bloom (2) increased population of fish due to lots of mortality of fish due to lack of oxygen V (4) increased population of aquatic organisms due to minerals T_{2} The highest DDT concentration in aquatic food chain shall occur in (1) seagull $T_{2} = -g Re^{2}M$ (4) phytoplankton $T_{2} = -g Re^{2}M$ (4) phytoplankton $T_{2} = -g Re^{2}M$ (5) eel. (6) phytoplankton $T_{2} = -g Re^{2}M$ (7) P.T.O.	/ identical molecules of single-stranded	
(2) HIV is unenveloped retrovirus. (3) HIV does not escape but attacks the acquired immune response. (4) HIV is enveloped virus containing one molecule of single-stranded RNA and one molecule of reverse transcriptase. (4) HIV is enveloped virus containing one molecule of reverse transcriptase. (5) Among the following edible fishes, which one is a marine fish having rich source of omega-3 fatty acids? (1) Mangur $g' = g' + $		
 (3) HIV does not escape but attacks the acquired immune response. (1) Allour Baliguo National Tark, Manipur 72 (1) (1) (1) Allour Baliguo National Tark, Manipur 72 (1) (1) (1) Allour Baliguo National Tark, Manipur 72 (1) (1) (1) (1) (1) (1) (1) (1) (1) (1)		
 (4) HIV is enveloped virus containing one molecule of single-stranded RNA and one molecule of reverse transcriptase. (1) Among the following edible fishes, which one is a marine fish having rich source of omega-3 fatty acids? (1) Mangur (2) Mrigala (3) Mackerel (4) Mystus (5) Michael (6) Michael (7) Simple and the following edible fishes, which one is a marine fish having rich source of omega-3 fatty acids? (1) Mangur (2) Mrigala (3) Mackerel (4) Mystus (5) Michael (6) Michael (7) Simple and the following edible fishes, which one is a marine fish having rich source of omega-3 fatty acids? (1) Mangur (2) Mrigala (3) Mackerel (4) Mystus (5) Simple and the following edible fishes, which one is a marine fish having rich source of omega-3 fatty acids? (1) Mangur (2) Mrigala (3) Mackerel (4) Mystus (5) Simple and the following edible fishes, which one is a marine fish having rich source of omega-3 fatty acids? (1) Mangur (2) Mrigala (3) Mackerel (4) Mystus (5) Simple and the following edible fishes, which one is a marine fish following edible fishes, which one is a marine fish having rich source of the lake due to algal bloom (1) Mangur (2) Called to the lake due to algal bloom (3) Mackerel (4) Mystus (5) Called to the lake due to algal bloom (6) fish due to lack of oxygen (7) The highest DDT concentration in aquatic food chain shall occur in the lake due to algal bloom (8) eel (9) eel (1) phytoplankton (1) phytoplankton (2) Called to the lake due to algal bloom (3) Mackerel (4) phytoplankton (5) Called to the lake due to algal bloom (6) the lake due to algal bloom (7) The highest	(3) HIV does not escape but attacks the	124. A lake which is rich in organic (aste may
molecule of single-stranded RNA and one molecule of reverse transcriptase? (1) Among the following edible fishes, which one is a marine fish having rich source of omega-3 fatty acids? (1) Mangur (2) Mrigala (3) Mackerel (4) Mystus j' = j' =		11-/
Among the following edible fishes, which one is a marine fish having rich source of omega-3 fatty acids? (1) Mangur (2) Mrigala (3) Mackerel (4) increased population of aquatic organisms due to minerals (4) increased population of aquatic organisms due to minerals (1) Mangur (2) Mrigala (3) Mackerel (4) Mystus $9'=5(1-2L_{Re})$ (4) increased population of aquatic food chain shall occur in (1) seagull (2) crabo (3) eel (4) phytoplankton $TC = -\frac{9}{Re^2M}$ (4) phytoplankton $TC = -\frac{9}{Re^2M}$ (2) Creabo (3) eel (4) phytoplankton $TC = -\frac{9}{2}(Re+M)$ (4) phytoplankton $TC = -\frac{9}{2}(Re+M)$ (5) eel (6) eel (6) eel (6) eel (7) eel (7) eel (8) eel (9) eel (molecule of single-stranded RNA and one	(2) increased population of fish due to lots of
one is a marine fish having rich source of omega-3 fatty acids? (1) Mangur (2) Mrigala (3) Mackerel (4) Mystus g' = g(1-2k) MD/E4 g' = g(1-2k) g' = g(1-2k)	\wedge	
omega-3 fatty acids? (1) Mangur $g' = \frac{g}{(+h_{R})}$ The highest DDT concentration in aquatic (2) Mrigala $g' = \frac{g}{(+h_{R})}$ The highest DDT concentration in aquatic food chain shall occur in (1) seagull $TC = \frac{g}{M_{em}}$ The highest DDT concentration in aquatic (2) Crabo $TC = \frac{g}{R}$ The highest DDT concentration in aquatic (3) Mackerel $g' = g(1 - 2h_{R})$ The highest DDT concentration in aquatic (4) Mystus $g' = g(1 - 2h_{R})$ The highest DDT concentration in aquatic (4) phytoplankton $TC = -\frac{g}{R}$ The highest DDT concentration in aquatic (4) phytoplankton $TC = -\frac{g}{R}$ The highest DDT concentration in aquatic (4) phytoplankton $TC = -\frac{g}{R}$ The highest DDT concentration in aquatic (5) Concentration in aquatic (6) Concentration in aquatic (1) seagull $TC = -\frac{g}{R}$ The highest DDT concentration in aquatic (2) Crabo $TC = -\frac{g}{R}$ The highest DDT concentration in aquatic (2) Crabo $TC = -\frac{g}{R}$ The highest DDT concentration in aquatic (3) Model $TC = -\frac{g}{R}$ The highest DDT concentration in aquatic (4) phytoplankton $TC = -\frac{g}{R}$ The highest DDT concentration in aquatic (4) phytoplankton $TC = -\frac{g}{R}$ The highest DDT concentration in aquatic (5) Concentration in aquatic (6) Concentration in aquatic (7) Concentration in aquatic (8) Concentration in aquatic (9) Concentration in aquatic (1) seagull $TC = -\frac{g}{R}$ The highest DDT concentration in aquatic (1) seagull $TC = -\frac{g}{R}$ The highest DDT concentration in aquatic (1) seagull $TC = -\frac{g}{R}$ The highest DDT concentration in aquatic (2) Cree + Cherner (2) C	one is a marine fish having rich source of	
(1) Mangur (2) Mrigala (3) Mackerel (4) Mystus	omega-3 fatty acids?	
(4) Mystus $9=9(1-2h)$ (4) phytoplankton $TC = -\frac{9}{2}Re^2M$ (4) mD/E4 $51=9(1-2h)=(Rt^2h)^2$ (1) (4) phytoplankton $TC = -\frac{9}{2}Re^2M$ [P.T.O.	(1) Mangur $q' = 9$	food chain shall occur in
(4) Mystus $9=9(1-2h)$ (4) phytoplankton $TC = -\frac{9}{2}Re^2M$ (4) mD/E4 $51=9(1-2h)=(Rt^2h)^2$ (1) (4) phytoplankton $TC = -\frac{9}{2}Re^2M$ [P.T.O.	(2) Mrigala U (+h)	(1) seaguil $TF = TP$
MD/E4 $\mathfrak{I} = \mathfrak{I} \left(1 - \frac{2h}{R} \right) = (Rtah) \mathfrak{I}_{\mathfrak{I}}$ [P.T.O.	(3) Mackerel a SI 21	20.200
	(4) Mystus $\int = \int \left(-\frac{2\pi}{R_e} \right)$	(4) phytoplankton TC = - g Re ² M
		2(Re+n)
$Te = -\frac{mRe^2}{2(Re+h)} \begin{pmatrix} R-2h \\ R \end{pmatrix} go$ $TE = -\frac{mgRe^2}{2(Re+h)}$ $Z(Re+h)$	9' - 9(1 - 20) = (k)	P.T.O.
2(Reth) (P) go Z(Reth)	To MOLZIN DUNE	TF= - mgfe ²
-INETHIN	$r = \frac{1}{2! Partial} \left(\frac{k - 2N}{2} \right) gc$	2(Re+h)
	-INETIN ()) U	



$$D(1 = D) - TDS \qquad Mark convert M + M - \pi$$


$$136. Which one of the following is incorrect for ideal solution?
$$(1) \Delta D_{mix} = 0 \cdot (1) \Delta D_{mix} =$$$$

1

f

148. Hot concentrated sulphuric acid is a 154. Jahn-Teller effect is not observed in high spin complexes of moderately strong oxidizing agent. Which of the following reactions does not show (1) d^8 (2) d^4 oxidizing behaviour? (1) $3S + 2H_2SO_4 \rightarrow 3SO_2 + 2H_2O$ (3) d^9 (4) d^7 (2) $C + 2H_2SO_4 \rightarrow CO_2 + 2SO_2 + 2H_2O_3$ (2) $CaF_2 + H_2SO_4 \rightarrow CaSO_4^7 + 2H_7^7$ 155. Which of the following can be used as (4) $Cu + 2H_2SO_4 \rightarrow CuSO_4 + SO_2 + 2H_2O_4$ the halide component for Friedel-Crafts 149. Which of the following pairs of d-orbitals will reaction? have electron density along the axes? (1) Bromobenzene (1) d_{xz}, d_{yz} (2) d 2, d 2 - 12 (2) Chloroethene (3) d_{xy} ; $d_{x^2-u^2}$ (3) Isopropyl chloride (4) d_{2}, d_{xz} (4) Chlorobenzene 150. The correct geometry and hybridization for XeF₄ are 56. In which of the following molecules, all (1) trigonal bipyramidal, sp^3d atoms are coplanar? (2) planar triangle, sp^3d^3 (2)(3) square planar, sp^3d^2 (4) octahedral, sp^3d^2 (4) -6 151. Among the following, which one is a wrong ISP3 & statement? (3)(1) $p\pi$ - $d\pi$ bonds are present in SO₂. (2) SeF₄ and CH_4 have same shape. (3) I_3^+ has bent geometry. 157. Which one of the following structures (4) PH5 and BiCl5 do not exist. represents nylon 6,6 polymer? 152. The correct increasing order of trans-effect H_2 H_2 H_2 H_2 H_1 H_2 H_2 of the following species is (1) $CN^- > C_6H_5^- > Br^- > NH_3$ (2) $Br^- > CN^- > NH_3 > C_6H_5^-$ (3) $CN^- > Br^- > C_6H_5^- > NH_3$ (4) $NH_3 > CN^- > Br^- > C_6H_5^-$ 153. Which one of the following statements related to lanthanons is incorrect? H-+CH2)6-NH (1) The basicity decreases as the ionic radius decreases from Pr to Lu. (2) All the lanthanons are much more reactive than aluminium. C^{H_2} H_2 C^{H_2} H_2 $H_$ (3) Ce(+4) solutions are widely used as oxidizing agent in volumetric analysis. (4) Europium shows +2 oxidation state. 16 JMD/E4 can-let 2 of court

58. In pyrrole



the electron density is maximum on (1) 3 and 4

- (2) 2 and 4
- (3) 2 and 5
- (4) 2 and 3

....

Which of the following compounds shall **not** produce propene by reaction with HBr followed by **e**limination or direct only elimination reaction?

(1)
$$H_3C - C - CH_2OH CH_2 - CH = CH_2$$

(1) $H_3C - C - CH_2OH CH_2 - CH = CH_2$
(3) $H_3C - C - CH_2Br$
(4) $H_2C - CH_2 - CH_2 - CH_2$
 H_2

160. Which one of the following nitro-compounds does not react with nitrous acid?

(1)
$$H_3C$$
 CH NO_2
 H_3C

 $\begin{array}{c} H_{3}C \\ H_{3}C \\ H_{3}C \\ H_{3}C \end{array}$

 $\begin{array}{c} (4) \quad H_3C \\ H_2 \end{array} \xrightarrow{C^2} NO_2 \end{array}$

JMD/E4

161. The central dogma of molecular genetics states that the genetic information flows from

(1) DNA \rightarrow Carbohydrates \rightarrow Proteins


(2) DNA \rightarrow RNA \rightarrow Proteins

- (3) DNA \rightarrow RNA \rightarrow Carbohydrates
- (4) Amino acids \rightarrow Proteins \rightarrow DNA
- 162. The correct corresponding order of names of four aldoses with configuration given below


respectively, is


- (1) D-threose, D-erythrose, L-threose, L-erythrose
- (2) L-erythrose, L-threose, D-erythrose, D-threose
- (3) D-erythrose, D-threose, L-erythrose, L-threose
- (4) L-erythrose, L-threose, L-erythrose, D-threose


163. In the given reaction




the product P is








[P.T.O.



164. A given nitrogen-containing aromatic 1 compound A reacts with Sn/HCl, followed by HNO_2 to give an unstable compound B. B, on treatment with phenol, forms a beautiful coloured compound C with the molecular formula $C_{12}H_{10}N_2O$. The structure of compound A is



165. Consider the reaction


 $CH_3CH_2CH_2Br + NaCN \rightarrow CH_3CH_2CH_2CN + NaBr$

This reaction will be the fastest in


- (1) methanol $\int \mathcal{I} = \int \mathcal{I}$
- (2) N, N'-dimethylformamide (DMF) $J_{MON} \simeq (N + \sqrt{12})^2$ (3) water $\equiv (\sqrt{N} + 1)^2$
- (4) ethanol = $n+1+2 \ln n$ $\int \min d (m-1)^2 = n+1-2 \ln n$


166. The **correct** structure of the product A formed in the reaction

given nitrogen-containing aromatic 167. Which among the given molecules can exhibit tautomerism?



- (1) Both I and III
- (2) Both I and II
- (3) Both II and III
- (4) III only





$$\frac{H_{2}(gas, 1 \text{ atmosphere})}{Pd/carbon, ethanol} A$$

$$\frac{Pmon-Jmin}{Iman+Jmin} = (n+1) + 2fn - (n+1) - fn^{169}, The compound that will react most readily with gaseous bromine has the formula with gaseous bromine has the formula of there is the formula of th$$