## HLAAC



Test Booklet Code

This Booklet contains 24 pages.



Do not open this Test Booklet until you are asked to do so.

## Read carefully the Instructions on the Back Cover of this Test Booklet.

## **Important Instructions :**

- The Answer Sheet is inside this Test Booklet. When you are directed to open the Test Booklet,
   take out the Answer Sheet and fill in the particulars on Side-1 and Side-2 carefully with blue/black ball point pen only.
- The test is of 3 hours duration and this Test Booklet contains 180 questions. Each question carries 4 marks. For each correct response, the candidate will get 4 marks. For each incorrect response, one mark will be deducted from the total scores. The maximum marks are 720.
- 3. Use Blue/Black Ball Point Pen only for writing particulars on this page/marking responses.
- 4. Rough work is to be done on the space provided for this purpose in the Test Booklet only.
- 5. On completion of the test, the candidate must hand over the Answer Sheet to the Invigilator before leaving the Room/Hall. The candidates are allowed to take away this Test Booklet with them.
- 6. The CODE for this Booklet is KK. Make sure that the CODE printed on Side-2 of the Answer Sheet is the same as that on this Test Booklet. In case of discrepancy, the candidate should immediately report the matter to the Invigilator for replacement of both the Test Booklet and the Answer Sheet.
- The candidates should ensure that the Answer Sheet is not folded. Do not make any stray
  marks on the Answer Sheet. Do not write your Roll No. anywhere else except in the specified
  space in the Test Booklet/Answer Sheet.
- 8. Use of white fluid for correction is not permissible on the Answer Sheet.

The kinetic energies of a planet in an elliptical orbit about the Sun, at positions A, B and C are K<sub>A</sub>, K<sub>B</sub> and K<sub>C</sub>, respectively. AC is the major axis and SB is perpendicular to AC at the position of the Sun S as shown in the figure. Then



- (1)  $K_A > K_B > K_C$ .
- (2) K<sub>B</sub> < K<sub>A</sub> < K<sub>C</sub>
- (3)  $K_A < K_B < K_C$
- $(4) \quad K_{\rm R} > K_{\rm A} > K_{\rm C}$
- 2. A solid sphere is in rolling motion. In rolling motion a body possesses translational kinetic energy  $(K_t)$  as well as rotational kinetic energy  $(K_r)$  simultaneously. The ratio  $K_t : (K_t + K_r)$  for the sphere is
  - (1) 5:7 ,
  - (2) 10:7
  - (3) 7:10
  - (4) 2:5

S

- 3. A solid sphere is rotating freely about its symmetry axis in free space. The radius of the sphere is increased keeping its mass same. Which of the following physical quantities would remain constant for the sphere ?
  - (1) Moment of inertia
  - (2) Rotational kinetic energy
  - (3) Angular velocity
  - (4) Angular momentum.
  - If the mass of the Sun were ten times smaller and the universal gravitational constant were ten times larger in magnitude, which of the following is **not** correct ?
    - Walking on the ground would become more difficult.
    - (2) Time period of a simple pendulum on the Earth would decrease.
    - (3) Raindrops will fall faster.

(4) 'g' on the Earth will not change.


HLAAC/KK/Page 2

A toy car with charge q moves on a frictionless horizontal plane surface under the influence of a uniform electric field  $\vec{E}$ . Due to the force q  $\vec{E}$ , its velocity increases from 0 to 6 m/s in one second duration. At that instant the direction of the field is reversed. The car continues to move for two more seconds under the influence of this field. The average velocity and the average speed of the toy car between 0 to 3 seconds are respectively

- (1) 1 m/s, 3 m/s
- (2) 1 m/s, 3.5 m/s
- (3) 2 m/s, 4 m/s

6.

- (4) 1.5 m/s, 3 m/s
- A block of mass m is placed on a smooth inclined wedge ABC of inclination  $\theta$  as shown in the figure. The wedge is given an acceleration 'a' towards the right. The relation between a and 0 for the block to remain stationary on the wedge is



(2) 
$$-7i - 8j - 4k$$
  
(3)  $-8i - 4i - 7k$ 

(4) -7i - 4j - 8k

A student measured the diameter of a small steel ball using a screw gauge of least count 0.001 cm. The main scale reading is 5 mm and 'zero' of circular scale division coincides with 25 divisions above the reference level. If screw gauge has a zero error of -0.004 cm, the correct diameter of the ball is


(1) 0.525 cm

- (2) 0.053 cm
   (3) 0.521 cm
- (3) 0·521 cm
   (4) 0·529 cm

4) 0.979 cm

SPACE FOR ROUGH WORK

The volume (V) of a monatomic gas varies with its temperature (T), as shown in the graph. The ratio of work done by the gas, to the heat absorbed by it, when it undergoes a change from state A to state B, is



10. The fundamental frequency in an open organ pipe is equal to the third harmonic of a closed organ pipe. If the length of the closed organ pipe is 20 cm, the length of the open organ pipe is

- (2) 12.5 cm
- (3) 13·2 cm
- (4) 16 cm
- (1) At what temperature will the rms speed of oxygen molecules become just sufficient for escaping from the Earth's atmosphere ? (Given :

Mass of oxygen molecule (m) =  $2.76 \times 10^{-26}$  kg

Boltzmann's constant  $k_{\rm B}$  = 1.38  $\times\,10^{-23}$  J  ${\rm K}^{-1})$ 

(1)  $8.360 \times 10^4 \text{ K}$ 

- (2)  $5.016 \times 10^4 \text{ K}$
- (3)  $2.508 \times 10^4 \text{ K}$
- (4)  $1.254 \times 10^4 \text{ K}$

The efficiency of an ideal heat engine working between the freezing point and boiling point of water, is 2734 3734

- (1) 20%
- (2) 6.25%
- (3) 26.8%
- (4) 12.5%


A carbon resistor of  $(47 \pm 4.7)$  k $\Omega$  is to be marked with rings of different colours for its identification. The colour code sequence will be

R2

- (1) Yellow Violet Orange Silver
- (2) Yellow Green Violet Gold
- (3) Violet Yellow Orange Silver
- (4) Green Orange Violet Gold
- 14. A set of 'n' equal resistors, of value 'R' each, are connected in <u>series</u> to a battery of emf 'E' and internal resistance 'R'. The current drawn is I. Now, the 'n' resistors are connected in parallel to the same battery. Then the current drawn from battery becomes 10 I. The value of 'n' is
  - (1) 11
  - (2) 20
  - (3) 10°
  - (4) 9

1.20

A battery consists of a variable number 'n' of identical cells (having internal resistance 'r' each) which are connected in series. The terminals of the battery are short-circuited and the current I is measured. Which of the graphs shows the correct relationship between I and n?



English

HLAAC/KK/Page 3 KT = 1 T = 1T

9.

16. An em wave is propagating in a medium with a velocity  $\overrightarrow{V} = V \hat{i}$ . The instantaneous oscillating electric field of this em wave is along +y axis. Then the direction of oscillating magnetic field of

- the em wave will be along
- (1) + z direction (2) y direction
- (3) -z direction
- (4) -x direction

17. The refractive index of the material of a prism is √2 and the angle of the prism is 30°. One of the two refracting surfaces of the prism is made a mirror inwards, by silver coating. A beam of monochromatic light entering the prism from the other face will retrace its path (after reflection from the silvered surface) if its angle of incidence on the prism is

- (1) 45°
- (2) 30°
- (3) 60°
- (4) zero

An object is placed at a distance of 40 cm from a concave mirror of focal length 15 cm. If the object is displaced through a distance of 20 cm towards the mirror, the displacement of the image will be

- (1) 36 cm away from the mirror.
- (2) 30 cm towards the mirror
- (3) 30 cm away from the mirror
- (4) 36 cm towards the mirror
- 19. The magnetic potential energy stored in a certain inductor is 25 mJ, when the current in the inductor is 60 mA. This inductor is of inductance
  - (1) 138·88 H
  - $(2) \quad 1.389 \ H$
  - (3) 0·138 H
  - (4) 13·89 H

HLAAC/KK/Page 4

An electron of mass m with an initial velocity  $\overrightarrow{V} = \overrightarrow{V_0} \stackrel{\circ}{i} (\overrightarrow{V_0} > 0)$  enters an electric field  $\overrightarrow{E} = -\overrightarrow{E_0} \stackrel{\circ}{i} (\overrightarrow{E_0} = \text{constant} > 0)$  at t = 0. If  $\lambda_0$  is its de-Broglie wavelength initially, then its de-Broglie wavelength at time t is


(1) 
$$\lambda_0 \left( 1 + \frac{eE_0}{mV_0} t \right)$$
  
(2)  $\lambda_0 t$   
(3)  $\frac{\lambda_0}{\left( 1 + \frac{eE_0}{mV_0} t \right)}$   
(4)  $\lambda_0$   
 $V = V_0 \lambda$   
 $V = V_0 \lambda$   
 $V = V_0 \lambda$   
 $F = eV_0 \lambda$   

- For a radioactive material, half-life is 10 minutes. If initially there are 600 number o nuclei, the time taken (in minutes) for the disintegration of 450 nuclei is
  - (1) 10
  - (2) 30
  - (3)= 20,
  - (4) 15
- 22. The ratio of kinetic energy to the total energy o an electron in a Bohr orbit of the hydrogen atom is
  - (1) 1:-1,
  - (2) 2:-1
  - (3) 1:1
  - (4) 1:-2
- 23. When the light of frequency  $2v_0$  (where  $v_0$  i threshold frequency), is incident on a meta plate, the maximum velocity of electrons emitter is  $v_1$ . When the frequency of the inciden radiation is increased to  $5v_0$ , the maximum velocity of electrons emitted from the same plat is  $v_2$ . The ratio of  $v_1$  to  $v_2$  is
  - (1) 1:4
  - (2) 4:1
  - (3) 1:2
  - (4) 2:1


SPACE FOR ROUGH WORK


- 24. Unpolarised light is incident from air on a plane surface of a material of refractive index 'μ'. At a particular angle of incidence 'i', it is found that the reflected and refracted rays are perpendicular to each other. Which of the following options is correct for this situation ?
  - (1) Reflected light is polarised with its electric
     vector perpendicular to the plane of
     incidence
    - (2)  $i = \sin^{-1}\left(\frac{1}{\mu}\right)$
    - (3) Reflected light is polarised with its electric vector parallel to the plane of incidence
    - $(4) \quad i = \tan^{-1} \left( \frac{1}{\mu} \right)$
- 25. An astronomical refracting telescope will have large angular magnification and high angular resolution, when it has an objective lens of
  - (1) large focal length and small diameter
  - (2) large focal length and large diameter
  - (3) small focal length and large diameter
  - (4) small focal length and small diameter
- 26. In Young's double slit experiment the separation d between the slits is 2 mm, the wavelength  $\lambda$  of the light used is 5896 Å and distance D between the screen and slits is 100 cm. It is found that the angular width of the fringes is 0.20°. To increase the fringe angular width to 0.21° (with same  $\lambda$ and D) the separation between the slits needs to be changed to
  - (1) 1.9 mm
  - (2) 2·1 mm
  - (3) 1·8 mm
  - (4) 1·7 mm

. In the circuit shown in the figure, the input voltage  $V_i$  is 20 V,  $V_{BE} = 0$  and  $V_{CE} = 0$ . The values of  $I_B$ ,  $I_C$  and  $\beta$  are given by



- (4) affects the overall V I characteristics of p-n junction
- 29. In the combination of the following gates the output Y can be written in terms of inputs A and B as






A tuning fork is used to produce resonance in a 34) glass tube. The length of the air column in this tube can be adjusted by a variable piston. At room temperature of 27°C two successive resonances are produced at 20 cm and 73 cm of column length. If the frequency of the tuning fork is 320 Hz, the velocity of sound in air at 27°C is

(1)339 m/s

80.


- (2)350 m/s
- 330 m/s (3)
- (4)300 m/s
- The electrostatic force between the metal plates 31. of an isolated parallel plate capacitor C having a charge Q and area A, is
  - distance linearly proportional to the (1)between the plates.
  - proportional to the square root of the (2)distance between the plates.
  - independent of the distance between the (3)plates. \*
  - inversely proportional to the distance (4)between the plates.
- An electron falls from rest through a vertical 36. (32) distance h in a uniform and vertically upward directed electric field E. The direction of electric field is now reversed, keeping its magnitude the same. A proton is allowed to fall from rest in it through the same vertical distance h. The time of fall of the electron, in comparison to the time of fall of the proton is
  - (1)5 times greater
  - (2)10 times greater
  - (3)smaller
  - (4)equal
- 33. A pendulum is hung from the roof of a sufficiently high building and is moving freely to and fro like a simple harmonic oscillator. The acceleration of the bob of the pendulum is 20 m/s<sup>2</sup> at a distance of 5 m from the mean position. The time period of oscillation is



- A metallic rod of mass per unit length 0.5 kg m<sup>-1</sup> is lying horizontally on a smooth inclined plane which makes an angle of 30° with the horizontal. The rod is not allowed to slide down by flowing a current through it when a magnetic field of induction 0.25 T is acting on it in the vertical direction. The current flowing in the rod to keep it stationary is A= 0.5
- 5.98 A (1)
- (2)14.76 A
- 7.14 A (3)

35.

- 11.32 A (4)
- Current sensitivity of a moving coil galvanometer is 5 div/mA and its voltage sensitivity (angular deflection per unit voltage applied) is 20 div/V. The resistance of the galvanometer is
- SI=5 SENA  $25 \Omega$ (1)250 Ω SY = NABXI SX1033 C X R (2)40 Q (3)(4)500 Ω
- A thin diamagnetic rod is placed vertically between the poles of an electromagnet. When the current in the electromagnet is switched on, then the diamagnetic rod is pushed up, out of the horizontal magnetic field. Hence the rod gains gravitational potential energy. The work required to do this comes from
- the magnetic field (1)
- the lattice structure of the material of the (2)rod
- (3)the current source
- the induced electric field due to the (4)changing magnetic field
- An inductor 20 mH, a capacitor 100 µF and a 37. resistor 50  $\Omega$  are connected in series across a source of emf,  $V = 10 \sin 314 t$ . The power loss in the circuit is
  - (1)0.43 W
  - (2)2.74 W
  - (3)0.79 W
  - (4)1.13 W



- 38. The power radiated by a black body is P and it radiates maximum energy at wavelength,  $\lambda_0$ . If the temperature of the black body is now changed so that it radiates maximum energy at wavelength  $\frac{3}{4}\lambda_0$ , the power radiated by it becomes nP. The value of n is
  - (1)  $\frac{4}{3}$ (2)  $\frac{256}{81}$ ,  $\lambda = P = T^{4}$ (3)  $\frac{3}{4}$ (4)  $\frac{81}{256}$
- 39. Two wires are made of the same material and have the same volume. The first wire has cross-sectional area A and the second wire has cross-sectional area 3A. If the length of the first wire is increased by Δl on applying a force F, how much force is needed to stretch the second wire by the same amount?

(1) 
$$6F + 7 = 72$$
  
(2)  $4F = 7$   
(3)  $9F = F = 74$   
(4)  $F = 72$   
(5)  $7 = FL = 74$   
(6)  $7 = FL = 74$   
(7)  $7 = FL = 74$   
(7)  $4 = FL = 74$   
(7)  $4 = FL = 74$   
(8)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 = 74$   
(9)  $7 =$ 


40. A small sphere of radius 'r' falls from rest in a viscous liquid. As a result, heat is produced due to viscous force. The rate of production of heat when the sphere attains its terminal velocity, is proportional to

- (2) r
- (3)  $r^{3}$
- (4) r<sup>4</sup>

41. A sample of 0.1 g of water at 100°C and normal pressure (1.013 × 10<sup>5</sup> Nm<sup>-2</sup>) requires 54 cal of heat energy to convert to steam at 100°C. If the volume of the steam produced is 167.1 cc, the change in internal energy of the sample, is

- (1) 208·7 J
- (2) 42.2 J
- (3) 104·3 J
- (4) 84·5 J

A body initially at rest and sliding along a frictionless track from a height h (as shown in the figure) just completes a vertical circle of diameter AB = D. The height h is equal to




Three objects, A : (a solid sphere), B : (a thin circular disk) and C : (a circular ring), each have the same mass M and radius R. They all spin with the same angular speed  $\omega$  about their own symmetry axes. The amounts of work (W) required to bring them to rest, would satisfy the relation

- (1)  $W_A > W_B > W_C$
- (2)  $W_B > W_A > W_C$
- (3)  $W_C > W_B > W_A$  (
- (4)  $W_A > W_C > W_B$
- A moving block having mass m, collides with another stationary block having mass 4m. The lighter block comes to rest after collision. When the initial velocity of the lighter block is v, then the value of coefficient of restitution (e) will be
  - (1) 0.25 •
  - (2) 0.8
  - (3) 0.5
  - (4) 0.4

Which one of the following statements is *incorrect*?

- Limiting value of static friction is directly proportional to normal reaction.
- (2) Frictional force opposes the relative motion.
- (3) Rolling friction is smaller than slidingfriction.
- (4) Coefficient of sliding friction has dimensions of length. •



					of the ions given i correct code :			I is treated with conc. $H_2SO_4$ . The evol- eous mixture is passed through KOH pello
			mn I		Column II		Wei	ight (in g) of the remaining product at S
	a.	Co <sup>3+</sup>		i	$\sqrt{8}$ B.M.		will	
	b.	Cr <sup>3+</sup>		/ti.	$\sqrt{35}$ B.M.		(1)	3.0
	c.	Fe <sup>3+</sup>	il.	iii.	$\sqrt{3}$ B.M.		(2)	2.8
	d.	Ni <sup>2+</sup>	10	iv.	$\sqrt{24}$ B.M.		(3) (4)	1·4 4·4
201	u.			v.	$\sqrt{15}$ B.M.	52.		difference between amylose and amyloped
			L.	1			is	unterence between anytose and anytoper
	245	a	ь	c	d		(1)	Amylose have $1 \rightarrow 4$ $\alpha$ -linkage a
	(1)	i	ii	iii	iv	1.1		$1 \to 6 \; \beta\text{-linkage}$
	(2)	iv	i	ii	iii		(2)	Amylopectin have $1 \rightarrow 4$ α-linkage a
	(3)	iv	v	ii	i •		(3)	$1 \rightarrow 6 \beta$ -linkage Amylopectin have $1 \rightarrow 4 \alpha$ -linkage a
0	(4)	iii	v	i	ii		(0)	$1 \rightarrow 6 \alpha$ -linkage .
47.)				(CO) <sub>5</sub> is			(4)	Amylose is made up of glucose a
	(1)		onuclea	r.		5		galactose
	(2)		clear			53.		arding cross-linked or network polyme
	(3) (4)		nuclear		<i>•</i>	11.4		ch of the following statements is <i>incorrect</i>
			clear		11 I	1.1	(1)	They are formed from bi- and tri-function monomers.
48,		type Cl <sub>2</sub> (en)		nerism s	hown by the complex	x	(2)	Examples are bakelite and melamine.
	(1)	1.1	-	n isomeri	sm		(3)	They contain covalent bonds betwee
	(2)			omerism			(1)	various linear polymer chains.
	(3)	Geon	netrical	isomeris	m		(4)	They contain strong covalent bonds in th polymer chains.
	(4)	Linka	age ison	nerism		54.	Nite	
49.	Whie	ch on	e of	the fol	lowing ions exhibits			ation of aniline in strong acidic medium a s m-nitroaniline because
					gnetism as well ?		(1)	In electrophilic substitution reaction
	(1)	$Cr_2O$	2- 7					amino group is meta directive.
	(2)	MnO	-			1	(2)	In absence of substituents nitro gro always goes to m-position. x
	(3)	$CrO_4^2$	-				(3)	In spite of substituents nitro group alwa goes to only m-position.
	(4)	MnO	2- 4 ×				(4)	In acidic (strong) medium aniline is prese as anilinium ion.
50.	The comj	geome olex [N	etry an li(CO) <sub>4</sub> ]	ld magn are	etic behaviour of the	55.	Whic natu	th of the following oxides is most acidic
	(1)	tetral	nedral g	geometry	and diamagnetic -		(1)	BeO -
	(2)	squar	e plana	r geomet	try and paramagnetic		(2)	BaO
	(3)	squar	e plana	r geomet	try and diamagnetic		(3)	MgO
	(4)	tetral	iedral g	eometry	and paramagnetic		(4)	CaO

L

56. In the reaction

$$\overset{OH}{\bigcirc} + CHCl_3 + NaOH \longrightarrow \overset{O^-Na^+}{\bigcirc} CHO$$

the electrophile involved is

- (1) formyl cation (CHO)
- (2) dichloromethyl anion (CHCl<sub>2</sub>)
- (3) dichloromethyl cation (CHCl<sub>2</sub>)
- (4) dichlorocarbene (:CCl<sub>2</sub>) ,
- 57. Carboxylic acids have higher boiling points than aldehydes, ketones and even alcohols of comparable molecular mass. It is due to their
  - (1) formation of carboxylate ion
  - (2) more extensive association of carboxylic acid via van der Waals force of attraction
  - (3) formation of intramolecular H-bonding
  - (4) formation of intermolecular H-bonding
- 58. Compound A, C<sub>8</sub>H<sub>10</sub>O, is found to react with NaOI (produced by reacting Y with NaOH) and yields a yellow precipitate with characteristic smell.

A and Y are respectively

(1) 
$$\bigcirc$$
 CH<sub>2</sub> – CH<sub>2</sub> – OH and I<sub>2</sub>

- (2)  $CH CH_3 \text{ and } I_2$ OH
- (3)  $H_3C \longrightarrow CH_2 OH \text{ and } I_2$

(4) 
$$CH_3 \longrightarrow OH \text{ and } I_2$$

HLAAC/KK/Page 9

SPACE FOR ROUGH WORK

- 59. The compound A on treatment with Na gives B, and with PCl<sub>5</sub> gives C. B and C react together to give diethyl ether. A, B and C are in the order
  - (1)  $C_2H_5OH$ ,  $C_2H_5Cl$ ,  $C_2H_5ONa$
  - $(2) \quad C_2H_5Cl, C_2H_6, C_2H_5OH$
  - (3)  $C_2H_5OH, C_2H_6, C_2H_5Cl$
  - (4)  $C_2H_5OH$ ,  $C_2H_5ONa$ ,  $C_2H_5Cl$

Which oxide of nitrogen is **not** a common pollutant introduced into the atmosphere both due to natural and human activity ?

- (1) NO<sub>2</sub>
- (2) N<sub>2</sub>O
- (3) N<sub>2</sub>O<sub>5</sub>
- (4) NO
- 61. Hydrocarbon (A) reacts with bromine by substitution to form an alkyl bromide which by Wurtz reaction is converted to gaseous hydrocarbon containing less than four carbon atoms. (A) is
  - (1)  $CH_2 = CH_2$
  - (2) CH<sub>3</sub> CH<sub>3</sub>
  - (3)  $CH \equiv CH$
  - (4) CH<sub>4</sub>,

$$C_7H_8 \xrightarrow{3 \operatorname{Cl}_2/\Delta} A \xrightarrow{\operatorname{Br}_2/\operatorname{Fe}} B \xrightarrow{\operatorname{Zn}/\operatorname{HCl}} C$$


The product 'C' is


- (1) o-bromotoluene
- (2) 3-bromo-2,4,6-trichlorotoluene
- (3) *m*-bromotoluene
- (4) p-bromotoluene.


English


60

63. Which of the following carbocations is expected to be most stable ?









64. Which of the following is correct with respect toI effect of the substituents ? (R = alkyl)

- (1)  $-NR_2 < -OR < -F$
- (2)  $-NH_2 > -OR > -F$
- $(3) NH_2 < -OR < -F$ .
- (4)  $-NR_2 > -OR > -F$
- 65. Which of the following molecules represents the order of hybridisation sp<sup>2</sup>, sp<sup>2</sup>, sp, sp from left to right atoms ?
  - (1)  $CH_2 = CH C \equiv CH$
  - (2)  $CH_2 = CH CH = CH_2$
  - (3)  $HC \equiv C C \equiv CH_{1}$
  - $(4) \quad CH_3 CH = CH CH_3$

HLAAC/KK/Page 10

Identify the major products P, Q and R in the following sequence of reactions :

- 67. Which of the following compounds can form zwitterion?
  - (1) Acetanilide
  - (2) Benzoic acid
  - (3) Aniline
  - (4) Glycine .

SPACE FOR ROUGH WORK

Engli

68.	Following solutions were prepared by mixing different volumes of NaOH and HCl of different concentrations :	72.	The bond dissociation energies of $X_2$ , $Y_2$ and $XY$ are in the ratio of $1: 0.5: 1$ . $\Delta H$ for the formation of XY is $-200$ kJ mol <sup>-1</sup> . The bond dissociation
	a. 60 mL $\frac{M}{10}$ HCl + 40 mL $\frac{M}{10}$ NaOH		energy of $X_2$ will be
	a. $60 \text{ mL} = \frac{10}{10} \text{ HCl} + 40 \text{ mL} = \frac{10}{10} \text{ NaOH}$		(1) $100 \text{ kJ mol}^{-1}$
	b. 55 mL $\frac{M}{10}$ HCl + 45 mL $\frac{M}{10}$ NaOH		(2) 800 kJ mol <sup>-1</sup> $\cdot$
	10 10		(3) $200 \text{ kJ mol}^{-1}$
	c. 75 mL $\frac{M}{5}$ HCl + 25 mL $\frac{M}{5}$ NaOH		(4) 400 kJ mol <sup>-1</sup>
	d. 100 mL $\frac{M}{10}$ HCl + 100 mL $\frac{M}{10}$ NaOH	73.	When initial concentration of the reactant is
	pH of which one of them will be equal to 1 ?		doubled, the half-life period of a zero order reaction
	(1) a		(1) is doubled ,
	(2) d		(2) is tripled
	(3) b (4) c.		(3) is halved
-	(4) Ce		(4) remains unchanged
(69.)	On which of the following properties does the coagulating power of an ion depend ?	74.	For the redox reaction
	(1) Size of the ion alone		$\operatorname{MnO}_4^- + \operatorname{C}_2\operatorname{O}_4^{2-} + \operatorname{H}^+ \longrightarrow \operatorname{Mn}^{2+} + \operatorname{CO}_2 + \operatorname{H}_2\operatorname{O}$
	(2) Both magnitude and sign of the charge on the ion		the correct coefficients of the reactants for the balanced equation are
	(3) The magnitude of the charge on the ion alone		$MnO_4^ C_2O_4^{2-}$ H <sup>+</sup>
	(4) The sign of charge on the ion alone		(1) 2 5 16
/79.	Given van der Waals constant for NH3, H2, O2		(2) 2 16 5
0	and $CO_2$ are respectively 4.17, 0.244, 1.36 and		(3) 16 5 2
	3.59, which one of the following gases is most		(4) 5 16 2
	easily liquefied ?	75.	Which one of the following conditions will favour
	(1) H <sub>2</sub>		maximum formation of the product in the reaction,
	(2) O <sub>2</sub>		$\mathbf{A}_{2}\left(\mathbf{g}\right)+\mathbf{B}_{2}\left(\mathbf{g}\right)\rightleftharpoons\mathbf{X}_{2}\left(\mathbf{g}\right)\  \   \Delta_{\mathbf{r}}\mathbf{H}=-\mathbf{X}\ \mathbf{kJ}\ ?$
	(3) NH <sub>3</sub>		(1) Low temperature and low pressure
	(4) CO <sub>2</sub>		(2) High temperature and high pressure
71.	The solubility of BaSO4 in water is		(3) Low temperature and high pressure .
	$2.42 \times 10^{-3}$ gL <sup>-1</sup> at 298 K. The value of its		(4) High temperature and low pressure
	solubility product (K <sub>sp</sub> ) will be	76.	The correction factor 'a' to the ideal gas equation
	(Given molar mass of $BaSO_4 = 233 \text{ g mol}^{-1}$ )		corresponds to
	(1) $1.08 \times 10^{-12} \text{ mol}^2 \text{ L}^{-2}$		(1) volume of the gas molecules
	(2) $1.08 \times 10^{-14} \text{ mol}^2 \text{ L}^{-2}$		(2) electric field present between the gas molecules
	(3) $1.08 \times 10^{-10} \text{ mol}^2 \text{ L}^{-2}$ ,		(3) density of the gas molecules
	(4) $1.08 \times 10^{-8} \text{ mol}^2 \text{ L}^{-2}$		(4) forces of attraction between the gas molecules.
		1	
HLA	AC/KK/Page 11 SPACE FOR I	ROUG	H WORK English

The correct difference between first-The correct order of N-compounds and 83. in its 77. second-order reactions is that decreasing order of oxidation states is HNO3, NO, NH4Cl, N2 the half-life of a first-order reaction does not (1)(1) depend on [A]<sub>0</sub>; the half-life of a HNO3, NH4Cl, NO, N2 (2)second-order reaction does depend on [A] HNO3, NO, N2, NH4Cl + (3)a first-order reaction can be catalyzed; a (2)second-order reaction cannot be catalyzed NH4Cl, N2, NO, HNO3 (4)the rate of a first-order reaction does not (3)depend on reactant concentrations; the rate 78. Which one of the following elements is unable to of a second-order reaction does depend on form  $MF_6^{3-}$  ion ? reactant concentrations (1)Al the rate of a first-order reaction does (4)depend on reactant concentrations; the rate (2)B . of a second-order reaction does not depend (3)Ga on reactant concentrations (4)In 84. Among CaH2, BeH2, BaH2, the order of ionic Considering Ellingham diagram, which of the character is following metals can be used to reduce alumina? (1)  $CaH_2 < BeH_2 < BaH_2$ (1)Zn (2)  $BeH_2 < BaH_2 < CaH_2$ (2)Mg (3)  $BeH_2 < CaH_2 < BaH_2 +$ (3)Fe  $BaH_2 < BeH_2 < CaH_2$ (4)(4)Cu In which case is the number of molecules of water 85) The correct order of atomic radii in group 13 80. maximum? elements is 0.18 g of water (1) B < Al < Ga < In < Tl0.00224 L of water vapours at 1 atm and (2)(2)B < Ga < Al < Tl < In 273 KB < Al < In < Ga < Tl (3)(3)18 mL of water  $10^{-3}$  mol of water (4) B < Ga < Al < In < Tl. (4)Which of the following statements is not true for /86. Consider the change in oxidation state of halogens? Bromine corresponding to different emf values as All are oxidizing agents. shown in the diagram below : (2)All but fluorine show positive oxidation  $BrO_4^- \xrightarrow{1.82 V} BrO_3^- \xrightarrow{1.5 V} HBrO$ states. (3) All form monobasic oxyacids.  $Br^{-} \leftarrow 1.0652 V Br_2 \leftarrow 1.595 V$ (4)Chlorine has the highest electron-gain enthalpy./ Then the species undergoing disproportionation 82. In the structure of ClF<sub>3</sub>, the number of lone pairs is of electrons on central atom 'Cl' is BrO<sub>4</sub> (1)(1)two (2) $Br_2$ (2)four (3)BrO. (3)one (4)three (4)HBrO HLAAC/KK/Page 12 SPACE FOR ROUGH WORK English

87. Consider the following species :

CN<sup>+</sup>, CN<sup>-</sup>, NO and CN

Which one of these will have the highest bond order ?

- (1) CN<sup>-</sup>\*
- (2)  $CN^+$
- (3) NO
- (4) CN

88. Which one is a *wrong* statement?

- An orbital is designated by three quantum numbers while an electron in an atom is designated by four quantum numbers.
- (2) The electronic configuration of N atom is

- (3) Total orbital angular momentum of electron in 's' orbital is equal to zero.
- (4) The value of m for  $d_{z^2}$  is zero.

89. Iron exhibits bcc structure at room temperature.
 Above 900°C, it transforms to fcc structure. The ratio of density of iron at room temperature to that at 900°C (assuming molar mass and atomic radii of iron remains constant with temperature) is

(1) 
$$\frac{4\sqrt{3}}{3\sqrt{2}}$$
  
(2)  $\frac{3\sqrt{3}}{3\sqrt{3}}$ 

$$(3) \quad \frac{\sqrt{3}}{\sqrt{3}}$$

$$(4) \quad \frac{1}{2}$$

- 90. Magnesium reacts with an element (X) to form an ionic compound. If the ground state electronic configuration of (X) is 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>3</sup>, the simplest ground for this compound is
  - (1) MgX<sub>2</sub>
  - (2) Mg<sub>2</sub>X
  - (3) Mg<sub>2</sub>X<sub>3</sub>
  - (4) Mg<sub>3</sub>X<sub>2</sub> •

HLAAC/KK/Page 13

- **91.** Pollen grains can be stored for several years in liquid nitrogen having a temperature of
  - (1)  $-80^{\circ}C$
  - (2) 196°C ·
  - $(3) 120^{\circ}C$
  - (4)  $-160^{\circ}C$
- 92. Oxygen is not produced during photosynthesis by(1) Nostoc
  - (2) Cycas
  - (3) Green sulphur bacteria.
  - (4) Chara
- **93.** What is the role of NAD<sup>+</sup> in cellular respiration?
  - (1) It functions as an electron carrier.
  - (2) It is a nucleotide source for ATP synthesis.
  - (3) It functions as an enzyme.
  - (4) It is the final electron acceptor for anaerobic respiration.
- **94.** Which of the following elements is responsible for maintaining turgor in cells ?
  - (1) Sodium
  - (2) Potassium.
  - (3) Magnesium
  - (4) Calcium
- **95.** Which one of the following plants shows a very close relationship with a species of moth, where none of the two can complete its life cycle without the other ?
  - (1) Yucca.
  - (2) Banana
  - (3) Hydrilla
  - (4) Viola
- In which of the following forms is iron absorbed by plants?
  - (1) Ferrous
  - (2) Free element
  - (3) Ferric
  - (4) Both ferric and ferrous

Double fertilization is

- Fusion of one male gamete with two polar nuclei
- (2) Fusion of two male gametes with one egg
- (3) Fusion of two male gametes of a pollen tube with two different eggs
- (4) Syngamy and triple fusion .

SPACE FOR ROUGH WORK

98. A 'new' variety of rice was patented by a foreign 10	<ul> <li>Niche is</li> <li>(1) the physical space where an organism lives</li> </ul>
98. A 'new' variety of rice was patented by deep company, though such varieties have been present in India for a long time. This is related to	(2) the range of temperature that are
(1) Sharbati Sonora	(3) all the biological factors in the organization
I Boin	
(3) Co-667	<ul> <li>(4) the functional role played by the organism</li> <li>where it lives •</li> </ul>
tion responsible for	<b>05.</b> Which of the following is a secondary pollutant ?
(99) In India, the organisation response assessing the safety of introducing genetically modified organisms for public use is and Industrial	(1) $CO_2$
(1) Council for Scientific and Industrial	(2) SO <sub>2</sub>
Description (CSIR)	(3) CO
Research Committee on General	(4) O <sub>3</sub> ,
at indiction (RCGM)	106. Natality refers to
Council of Medical Research (Terrate	T' il noto f
<ul> <li>(3) Indian Council of Appraisal Committee</li> <li>(4) Genetic Engineering Appraisal Committee</li> </ul>	<ol> <li>Birth rate</li> <li>Number of individuals leaving the habitat</li> </ol>
(GEAC)	
100. Which of the following is commonly used as a	<ul><li>(3) Death rate</li><li>(4) Number of individuals entering a habitat</li></ul>
vector for introducing a	
lymphocytes ?	107. World Ozone Day is celebrated on
(1) Ti plasmid	(1) 21 <sup>st</sup> April
(2) $\lambda$ phage	the success
(3) Retrovirus.	
(4) pBR 322	(3) 5 <sup>th</sup> June
a stinotional companies	e (4) 22 <sup>nd</sup> April.
101. Use of bioresources by multinational companies of and organisations without authorisation from the concerned country and its people is called	<ul><li>(1) 108. What type of ecological pyramid would obtained with the following data?</li></ul>
(1) Biopiracy "	Secondary consumer : 120 g
(2) Biodegradation	Primary consumer : 60 g
(3) Bio-infringement	Primary producer : 10 g
(4) Bioexploitation	
102. The correct order of steps in Polymerase Cha	(1) Pyramid of energy (2) Upright pyramid of numbers
	a fiomass .
Appending Extension, Denaturation	d of biomass
Departuration, Extension, Annealing	(4) Upright pyramid of official and a second
<ul> <li>(2) Denaturation, annealing</li> <li>(3) Extension, Denaturation, Annealing</li> </ul>	109. In stratosphere, which of the following elem
<ul> <li>(3) Extension, Denaturation, Annealing, Extension.</li> <li>(4) Denaturation, Annealing, Extension.</li> </ul>	
The is the correct match :	release of molecular oxygen
P Pagesive parent - Dinystia	
TIL Morgan - Transuterio	
- Nucleic acto	
(3)Ribozyme-Nuclei(4)G. Mendel-Transforma	
(4) G. Mendel SPACE	OR ROUGH WORK

HLAAC/KK/Page 14

-

	matched ?		Secondary xylem and phloem in dicot stem are produced by
	(1) ABO blood grouping : Co-dominance/		(1) Vascular cambium-
	(2) XO type sex : Grasshopper		(2) Phellogen
	determination		(3) Apical meristems
	(3) Starch synthesis in pea : Multiple alleles		(4) Axillary meristems
	(4) T.H. Morgan : Linkage	118.	Plants having little or no secondary growth are
111.	Select the <i>correct</i> statement :		(1) Deciduous angiosperms
	<ol> <li>Punnett square was developed by a British scientist.</li> </ol>		(2) Conifers
	(2) Spliceosomes take part in translation.		(3) Grasses
	<ul><li>(3) Franklin Stahl coined the term "linkage".</li></ul>		(4) Cycads (4)
	<ul><li>(4) Transduction was discovered by S. Altman.</li></ul>	119.	Sweet potato is a modified
GP			(1) Adventitious root
12)	The experimental proof for semiconservative replication of DNA was first shown in a		(2) Tap root •
	(1) Bacterium.		(3) Stem
	(2) Plant		(4) Rhizome
	(3) Fungus	120.	Pneumatophores occur in
	(4) Virus		(1) Free-floating hydrophytes
113.	Select the <i>correct</i> match :		<ul><li>(2) Carnivorous plants</li></ul>
	(1) Alfred Hershey and – TMV		(3) Halophytes.
	Martha Chase		(4) Submerged hydrophytes
	(2) Matthew Meselson – Pisum sativum	101	
	and F. Stahl	121.	Casparian strips occur in
	(3) Alec Jeffreys – Streptococcus		(1) Pericycle
	. pneumoniae		(2) Cortex,
	(4) Francois Jacob and $-Lac$ operon		(3) Epidermis
~	Jacques Monod		(4) Endodermis,
14.	Offsets are produced by	122.	Which of the following statements is <i>correct</i> ?
9	<ol> <li>Mitotic divisions</li> <li>Parthenocarpy.</li> </ol>		(1) Selaginella is heterosporous, while Salvinia
	(3) Meiotic divisions	1	<ul> <li>is homosporous. ×</li> <li>(2) Horsetails are gymnosperms. ×</li> </ul>
	(4) Parthenogenesis		<ul><li>(3) Ovules are not enclosed by ovary wall in</li></ul>
115.	Which of the following flowers only once in its life-time ?		gymnosperms. *
	(1) Jackfruit ·		(4) Stems are usually unbranched in both
	(2) Mango	1.0	Cycas and Cedrus.
	(3) Bamboo species,	123	Select the <i>wrong</i> statement :
	(4) Papaya		(1) Mushrooms belong to Basidiomycetes.
116.	Which of the following has proved helpful in preserving pollen as fossils ?		(2) Pseudopodia are locomotory and feeding structures in Sporozoans.
	(1) Cellulosic intine		<ul><li>(3) Cell wall is present in members of Fungi</li></ul>
	(2) Oil content		and Plantae.
	(3) Pollenkitt		(4) Mitochondria are the powerhouse of the cell
	(4) Sporopollenin •		in all kingdoms except Monera.

124.	Mat Colu	ch the items umn II and s	given select	in Colum the cor	nn I with those in rect option given	120.	sug	two functional groups characteristic of ars are
	belo						(1)	carbonyl and methyl
		Column I		Column	II		(2)	carbonyl and phosphate
	a.	Herbarium	i	Itisan	lace having a		(3)	hydroxyl and methyl
	a.	Tierbar fuin	1.		n of preserved	-	(4)	carbonyl and hydroxyl •
		1			nd animals.	(129.	Whi	ich of the following is <i>not</i> a product of light
	b.	Key *	ii.	· · · · · · · · · · · · · · · · · · ·	at enumerates	~	read	ction of photosynthesis ?
	D.	Rey	п.	an move on	ically all the		(1)	NADH
		1			found in an area		(2)	NADPH •
		)	1		ef description		(3)	ATP
					dentification.		(4)	Oxygen
	c.	Museum	liii.		e where dried and	130.	Whi	ich among the following is <i>not</i> a prokaryote ?
	с.	museum	m,		plant specimens		(1)	Mycobacterium
					d on sheets are		(2)	Nostoc
				kept.	a on oncoto are		(3)	Saccharomyces.
	d.	Catalogue	iv.		et containing a list	1.1	(4)	Oscillatoria
		enter Bar			cters and their	131.	Stor	matal movement is <i>not</i> affected by
				alternat	es which are		(1)	Light
				helpful	in identification of		(2)	$O_2$ concentration,
				various	the second se		(3)	Temperature
		a b	с	d	÷		(4)	$\rm CO_2$ concentration
	(1)	iii ii	i	iv		132.	The	Golgi complex participates in
	(2)	ii iv	iii	i			(1)	Formation of secretory vesicles.
	(3)	i iv	iii	ii			(2)	Respiration in bacteria
	(4)	iii iv	i	ii .			(3)	Fatty acid breakdown
		1.0.00			G 1		(4)	Activation of amino acid
125.					neiosis, spores are	133.	Whi	ch of the following is true for nucleolus?
	1.	luced exogene		in			(1)	It is a membrane-bound structure.
	(1)	Alternaria 🛪 Agaricus 🖇	C				(2)	It takes part in spindle formation.
	(2)	Neurospora	A				(3)	Larger nucleoli are present in dividing cells.
	(4)	Saccharomy					(4)	It is a site for active ribosomal RNA
	1.17	Succitaronty	icco y					synthesis
126.	Win	ged pollen gra	ains a	re presei	nt in	134.	The	stage during which separation of the paired
	(1)	Cycas						nologous chromosomes begins is
	(2)	Mango					(1)	Diplotene
	(3)	Mustard					(2)	Diakinesis .
	(4)	Pinus					(3)	Pachytene
197	Whi	ab one is tom	n alu	motohod	2		(4)	Zygotene
14.	(1)	ch one is <b>wro</b> Biflagellate			Brown algae	135.	Stor	nata in grass leaf are
	(2)	Gemma cup		ores -	Marchantia /		(1)	Kidney shaped
	(2)	Uniflagellat		netes -	Polysiphonia	m	(2)	Rectangular
	(4)	Unicellular			Chlorella		(3)	Dumb-bell shaped,
	1.41	omoonului	Bul				(4)	Barrel shaped

136.	Niss	l bodies are mainly composed of	142.	All o	f the f	followin	g are pa	art of	f an operon <i>except</i>
	(1)	DNA and RNA		(1)		tural g			
	(2)	Nucleic acids and SER		(2)		nhancer			
	(3)	Proteins and lipids		(3)	an oj	perator			4
	(4)	Free ribosomes and RER		(4)	a pro	moter			
137.	Whie (1) (2)	ch of these statements is <i>incorrect</i> ? Glycolysis occurs in cytosol. Glycolysis operates as long as it is supplied	143.	X	chrom rited	osomes by		ch	ndition on one of her romosome can be $\sim \times \times$
	(3) (4)	with NAD that can pick up hydrogen atoms. Enzymes of TCA cycle are present in mitochondrial matrix. Oxidative phosphorylation takes place in outer mitochondrial membrane.		<ol> <li>(1)</li> <li>(2)</li> <li>(3)</li> <li>(4)</li> </ol>	Only Only	daught	hildren ærs nd daug	hter	$\frac{x^{c} \times x^{c} \gamma}{x^{c}} \times \frac{x^{c} \gamma}{x^{c}} \times \frac{x^{c}}{x^{c}}$
138.		ch of the following terms describe human ition?		evol	ution		go de V	ries	, the mechanism of
	(1)	Thecodont, Diphyodont, Heterodont -		(1) (2)			variation	ne	
	(2)	Pleurodont, Monophyodont, Homodont 🛪		(2)			p mutat		
	(3)	Thecodont, Diphyodont, Homodont		(4)	1000	or muta	1. A.	10115	
	(4)	Pleurodont, Diphyodont, Heterodont 🖍	0.01						1
(139.)	(1) (2) (3) (4)	ct the <i>incorrect</i> match : Allosomes – Sex chromosomes Submetacentric – L-shaped chromososmes chromosomes Lampbrush – Diplotene bivalents chromosomes Polytene – Oocytes of amphibians		stra	nd of nence UGC ACC AGG	a gene.	What w canscrib GCAT CGAU GCAU	vill t	nce from the coding be the corresponding nRNA ?
140.)	Whi	chromosomes ch of the following events does <b>not</b> occur in ch endoplasmic reticulum? $f \downarrow \varsigma$			ımn I		-		umn I with those in orrect option given
-	(1)	Protein glycosylation			Colu	mn I			Column II
	(2)	Cleavage of signal peptide		a.	Proli	iferative	Phase	i.	Breakdown of
	(3)	Protein folding							endometrial lining
	(4)	Phospholipid synthesis		b.	Soor	etory Pl	1950	ji ji	Follicular Phase
141.	Man	y ribosomes may associate with a single				struatio			Luteal Phase
	mR! sim	NA to form multiple copies of a polypeptide ultaneously. Such strings of ribosomes are ned as		c.	a i	b iii	c ii		Lawai i nast
	(1)	Polyhedral bodies							
	(2)	Plastidome		(2)	ii 	iii 	i'.		
	(3)	Polysome .		(3)	iii	ii	i		
	(4)	Nucleosome		(4)	iii	i	ii		

.

	intera	n one actions is	s widel	y use	d in me	dical science for		diver	ng the following sets of examples for rgent evolution, select the <i>incorrect</i> option : Heart of bat, man and cheetah
		roductio		101011	CS :			(2)	Brain of bat, man and cheetah
1.1		Mutuali						(3)	Forelimbs of man, bat and cheetah
		Parasiti							Eye of octopus, bat and man .
	(Q)	Commen		1				(4)	
	(4)	Amensa	lism .				153.	Whie	ch of the following is not an autoimmune
148.	All o	of the f	ollowin	g ar	e inclu	ded in 'Ex-situ			ase ?
1 10.	conse	ervation'	except					1.1	Rheumatoid arthritis Alzheimer's disease. Ww acefylcholin
	(1)	Sacred (	groves	•				(2)	Alzheimer's disease, where the 10
	(2)	Botanic	al gard	ens "				(3)	Psoriasis
	(3)	Wildlife	safari	parks	3			(4)	Vitiligo
	(4)	Seed ba					154.		which disease does mosquito transmitted nogen cause chronic inflammation of
149.	Matc	h the ite	ems giv	en in	Colum	n I with those in		lym	phatic vessels ?
			and sele	ect th	e corr	ect option given		(1)	Ascariasis
	below				Co	lumn II		(2)	Ringworm disease
		Column		Se		-B radiation		(3)	Elephantiasis.
	a.	Eutrop		1. I		orestation		(4)	Amoebiasis
	b.	Sanitar					155	Con	version of milk to curd improves it
	c.	Snow b	lindnes	S	iii. Nu		155.	nut	ritional value by increasing the amount of
						ichment		(1)	Vitamin A
	d.	Jhum c		ion		ste disposal		(2)	Vitamin B <sub>12</sub> .
			b	c	d				Vitamin D
	(1)	1.1	iii	iv	ii			(3) (4)	Vitamin E
	(2)		iv	i	ii e				
	(3) (4)	100	i ii	iii iv	iv iii		156.	The of n	e similarity of bone structure in the forelimb nany vertebrates is an example of
					of a con	ntm		(1)	Analogy
150.		growing						(2)	Convergent evolution
	(1)				ndividu	are less than the		(3)	Homology -
	101		-		ind	pre-reproductive		(4)	Adaptive radiation
	(2)		luals ar	e equ	al in nu	imber.	157.	. Which of the following characteristics represent Inheritance of blood groups' in humans ?	
	(3)	pre-rep	product	ive in	dividua dividua	als are more than ls.•		a.	Dominance
	(4)					als are less than		b.	Co-dominance /
	(4)	the rep	product	ive in	dividua	lls.		c.	Multiple allele 🗸
		0.000						d.	Incomplete dominance
151.	Whi	ch part	of pop	py pla	ant is u	used to obtain the		e.	Polygenic inheritance,
	1.1	g "Smac						(1)	a, b and c
	(1)	Latex						(2)	b, d and e
	(2)	Roots						(3)	b, c and e .
	(3)	Flower						(4)	a, c and e
	(4)	Leave	s	-			1		

.

and the second second

158.	Horm	ones secreted by the placenta to maintain	162.		h of the following is	s an	amino acid derived
	pregn	nancy are					
	(1)	hCG, hPL, estrogens, relaxin, oxytocin		(1)	Ecdysone		
	(2)	hCG, hPL, progestogens, estrogens -		(2)	Estradiol		
	(3)	hCG, hPL, progestogens, prolactin		(3)	Epinephrine •		
		hCG, progestogens; estrogens, glucocorticoids >		(4)	Estriol		
159.	The c	contraceptive 'SAHELI'	163.	Whi ince	ch of the following prrectly paired with	stru its f	actures or regions is function ?
	(1)	increases the concentration of estrogen and prevents ovulation in females.		(1)	Limbic system	:	consists of fibre tracts that
	(2)	is an IUD.					interconnect different regions of
	(3)	blocks estrogen receptors in the uterus, preventing eggs from getting implanted.					brain; controls movement.
	(4)	is a post-coital contraceptive.		(2)	Hypothalamus	:	production of releasing hormones
160.	The from	amnion of mammalian embryo is derived					and regulation of temperature, hunger and thirst.
	(1)	endoderm and mesoderm		(0)	Medulla oblongata		controls respiration
	(2)	mesoderm and trophoblast		(3)	Medulla obioligata		and cardiovascular
	(3)	ectoderm and mesoderm ,					reflexes.
	(4)	ectoderm and endoderm		(4)	Corpus callosum	:	band of fibers connecting left and
	The	rmiation is					right cerebral hemispheres.
Spar	(1)	In spermiogenesis spermatozoa are formed, while in spermiation spermatids are	101		e transparent lens in place by	the	e human eye is held in
		formed.		(1)	ligaments attache	d to	the iris
	(2)		E I	(2)	smooth muscles at	tacl	hed to the iris
		cells are released into the cavity o seminiferous tubules, while in spermiation		(3)			
		spermatozoa are formed.»		(4)	smooth muscles a	ttac	hed to the ciliary body
	(3)	while in spermiation spermatozoa are		5. Wl	nificant role in osteo	por	
		formed.		(1)			
	(4)	In spermiogenesis spermatozoa are formed while in spermiation spermatozoa ar	e	(2)			
		released from sertoli cells into the cavity of	f	(3)			
		seminiferous tubules.		(4)	Parathyroid horn	one	and Prolactin >

HLAAC/KK/Page 19

· .

÷.

1.2 ÷

1

	(4)	iv	iii	ii	i			filament.
	(3)	iii	ii	i	iv		121	the myosin cross bridges and the
	(2)	î	iv	ii	iii		(4)	prevents the formation of bonds bet
	(1)	iii	i	iv	ii •			active sites on actin for myosin.
		a	b	с	d		(3)	binds to troponin to remove the maski
	d.	Resid	lual vol	lume	iv. 1000 – 1100 mL		5.M/	filament.
	с.	volur	ne	Reserve	'iii. 500 – 550 mL		(2)	it. detaches the myosin head from the
		volur	ne	/	\		(1)	activates the myosin ATPase by bindi
	b.			Reserve	ii. 1100 – 1200 mL	112.		rium is important in skeletal m traction because it
	a.		volum	e	i 2500 – 3000 mL	179	Cale	
	below	w : Colui	mn I		Column II		(4)	Emphysema
	Colu	mn I			correct option given		(3)	Anthracis
168.	Mate	h the	items		Column I with those in		(2)	Botulism
	(4)	ii	i	iii			(1)	Silicosis 3
	(3)	iii	i	ii .			resp	piratory disorder ?
	(1)	i	ii	iii		171.	Whi	ich of the following is an occupat
	(1)	a i	b iii	c ii			(4)	ii iii i.
			5		ventricle		(3)	iii ii i
	· ·				atrium and right	10	(2)	i iii ii
	c.	Semi	ilunary	valve hi			(1)	i ii iii
					ventricle and pulmonary artery			a b c
	b.	Bicu	spid va	lve \ii.			C.	Albumin iii. Defence mechan
	a.	Trict	ispid v	alve i.	and left ventricle	I -	b.	Globulin ii. Blood clotting
			mn I	due 1	Column II Between left atrium		a.	Fibrinogen i. Osmotic balanc
	belo	w :			- to			Column I Column II
167.					Column I with those in e correct option given		belo	
105				on of bron				umn II and select the <i>correct</i> option
	(4)	Decr	eased	respira		170.	Mat	tch the items given in Column I with th
	(3)			on of k surface,	oronchioles; Decreased		(4)	Parietal cells,
		Infla	mmati	on of bron	nchioles		(3)	Chief cells
	(2)			surface respira	tory surface;		(2)	Goblet cells
	(1)				bronchioles; Increased		(1)	Mucous cells
				pectively		M	1.11	p in erythropoiesis ?
	repr	esents	s the ]	lung con	ng options correctly litions in asthma and	1-1	/	ich of the following gastric cells ind

173.	Match the items given in Column I with those in Column II and select the <i>correct</i> option given below :					a mala analmanah from a famala analmanah 2
4	belo	ow∶				
		Colur	mn I		Column II	
	a.	Glyco	osuria	i.	Accumulation of uric	(3) Presence of a boat shaped sternum on the 9 <sup>th</sup> abdominal segment
					acid in joints	(4) Presence of anal cerci
	b.	Gout		ii.	Mass of crystallised salts within the kidney	
	c,	Rena	l calcul	į iii.	Inflammation in	characterized by crop and gizzard in its digestive system.
					glomeruli	(1) Reptilia
	d.	Glom neph	erular ritis	\iv.	Presence of glucose in urine	(2) Aves.
						(3) Amphibia
		а	b	с	d	(4) Osteichthyes
	(1)	ï	ii	iii	iv	177. Which one of these animals is not a
	(2)	ii	iii	i	iv	homeotherm?
	(3)	iii	ii	iv	i -	(1) Chelone
	(4)	iv	i	ii -	iii •	(2) Camelus
174	Mat	ch the	itome (	rivon i	n Column I with those i	· ·
174.					he <i>correct</i> option give	
	bela		and 5	ciece i	are correct option give	(4) Psittacula ,
	Dere	Colur	nn I		Column II	.178. Which of the following organisms are known as chief producers in the oceans?
		(Fund	ction)		(Part of Excretory	(1) Diatoms
					System)	(2) Cyanobacteria
	a.	Ultra	filtratio	op.	i. Henle's loop	(3) Dinoflagellates,
	b.	Conce	entratio	mX	ii. Ureter	(4) Euglenoids
	c.		sport of		iii. Urinary bladder	179. Which of the following animals does <u>not</u> undergo metamorphosis?
	- 6	urine			/ ,	(1) Tunicate
	d.	Stora	ge of u	rine	iv. Malpighian	(2), Moth
	*				corpuscle	(3) Earthworm,
					v. Proximal convoluted tubule	(4) Starfish
						180. Ciliates differ from all other protozoans in
		а	b	с	d	(1) having a contractile vacuole for removing
	(1)	iv	i	jii	iii •	excess water
	(2)	v	iv	i	ii	(2) using pseudopodia for capturing prey $\rho_{\infty}$
	(3)	iv	v	ii	iii	(3) using flagella for locomotion $\sqrt{2}$
	(4)	v	iv	i,	iii +	(4) having two types of nuclei
111.0	0.000	V/Daga	04		SPACE FOI	ROUGH WORK English

HLAAC/KK/Page 21

SPACE FOR ROUGH WORK

BOOKLET CODE - HLAAC (KK)

BOOKLET CODE - HLA			
Q.No. Answer	Q.No. Answer	Q.No. Answer	Q.No. Answer
1. (1)	46. (3)	91. (2)	136. (4)
2. (1)	47. (1)	92. (3)	137. (4)
3. (4)	48. (3)	93. (1)	138. (1)
4. (4)	49. (4)	94. (2)	139. (4)
5. (1)	50. (1)	95. (1)	140. (4)
6. (4)	51. (2)	96. (3)	141. (3)
7. (4)	52. (3)	97. (4)	142. (2)
8. (4)	53. (4)	98. (4)	143. (4)
9. (3)	54. (4)	99. (4)	144. (1)
10. (3)	55. (1)	100. (3)	145. (3)
11. (1)	56. (4)	101. (1)	146. (2)
12. (3)	57. (4)	102. (4)	147. (4)
13. (1)	58. (2)	103. (3)	148. (1)
14. (3)	59. (4)	104. (4)	149. (2)
15. (3)	60. (3)	105. (4)	150. (3)
16. (1)	61. (4)	106. (1)	151. (1)
17. (1)	62. (3)	107. (2)	152. (4)
18. (1)	63. (2)	108. (3)	153. (2)
19. (4)	64. (3)*	109. (1)	154. (3)
20. (3)	65. (1)	110. (3)	155. (2)
21. (3)	66. (4)	111. (1)	156. (3)
22. (1)	67. (4)	112. (1)	157. (1)
23. (3)	68. (4)	113. (4)	158. (2)
24. (1)	69. (2)	114. (1)	159. (3)
25. (2)	70. (3)	115. (3)	160. (3)
26. (1)	71. (3)	116. (4)	161. (4)
27. (4)	72. (2)	117. (1)	162. (3)
28. (4)	73. (1)	118. (3)	163. (1)
29. (1)	74. (1)	119. (1)	164. (3)
30. (1)	75. (3)	120. (3)	165. (2)
31. (3)	76. (4)	121. (4)	166. (3)
32. (3)	77. (3)	122. (3)	167. (3)
33. (1)	78. (2)	123. (2)	168. (1)
34. (4)	79. (2)	124. (4)	169. (4)
35. (2)	80. (4)	125. (2)	170. (4)
36. (3)	81. (2)	126. (4)	171. (1)
37. (3)	82. (1)	127. (3)	172. (3)
38. (2)	83. (1)	128. (4)	173. (4)
39. (3)	84. (3)	129. (1)	174. (1)
40. (2)	85. (3)	130. (3)	175. (1)
41. (1)	86. (4)	131. (2)	176. (2)
42. (4)	87. (1)	132. (1)	177. (1)
43. (3)	88. (2)	133. (4)	178. (1)
44. (1)	89. (2)	134. (1)	179. (3)
45. (4)	90. (4)	135. (3)	180. (4)
		100. (0)	